

Calculation of radiation reaction effect on orbital parameters in Kerr spacetime

Norichika Sago^{1,*}, and Ryuichi Fujita²

¹*Faculty of Arts and Science, Kyushu University, Fukuoka 819-0395, Japan*

²*CENTRA, Departamento de Física, Instituto Superior Técnico, Universidade de Lisboa, Avenida Rovisco Pais 1, Portugal*

*E-mail: sago@artsci.kyushu-u.ac.jp

.....
We calculate the secular changes of the orbital parameters of a point particle orbiting a Kerr black hole, due to the gravitational radiation reaction. For this purpose, we use the post-Newtonian (PN) approximation in the first order black hole perturbation theory, with the expansion with respect to the orbital eccentricity. In this work, the calculation is done up to the fourth post-Newtonian (4PN) order and to the sixth order of the eccentricity, including the effect of the absorption of gravitational waves by the black hole. We confirm that, in the Kerr case, the effect of the absorption appears at the 2.5PN order beyond the leading order in the secular change of the particle's energy and may induce a superradiance, as known previously for circular orbits. In addition, we find that the superradiance may be suppressed when the orbital plane inclines with respect to the equatorial plane of the central black hole. We also investigate the accuracy of the 4PN formulae by comparing to numerical results. If we require that the relative errors in the 4PN formulae are less than 10^{-5} , the parameter region to satisfy the condition will be $p \gtrsim 50$ for $e = 0.1$, $p \gtrsim 80$ for $e = 0.4$, and $p \gtrsim 120$ for $e = 0.7$ almost irrespective of the inclination angle nor the spin of the black hole, where p and e are the semi-latus rectum and the eccentricity of the orbit. The region can further be extended using an exponential resummation method to $p \gtrsim 40$ for $e = 0.1$, $p \gtrsim 60$ for $e = 0.4$, and $p \gtrsim 100$ for $e = 0.7$. Although we still need the higher order calculations of the PN approximation and the expansion with respect to the orbital eccentricity to apply for data analysis of gravitational waves, the results in this paper would be an important improvement from the previous work at the 2.5PN order, especially for large p region.
.....

Subject Index E01, E02, E31, E36

1. Introduction

The gravitational two-body problem is a fundamental issue in general relativity. This also attracts great interest in gravitational wave physics because binary inspirals are promising sources of gravitational waves which are expected to be detected directly by ongoing gravitational wave observatories in the world. Understanding the dynamics of binary system is required to predict the emitted gravitational waveforms accurately for efficient searches of the signal in observed data.

One of major approaches for this purpose is the gravitational self-force (GSF) picture in the black hole perturbation theory. In this picture, a binary is regarded as a point mass orbiting a black hole and the dynamics can be described by the equation of motion of the mass including the effect of the interaction with the self-field, that is, the GSF. After the formal expression of the GSF was presented by Mino, Sasaki and Tanaka [1] and Quinn and

Wald [2], a lot of efforts have been devoted to develop practical formulations and methods to calculate the GSF (for example, refer to [3] for the formulation of GSF, [4, 5] for the recent progress in practical calculations of GSF).

Although a lot of progress has been made, however, it is still challenging to calculate the GSF directly for general orbits, especially in Kerr spacetime. Practical calculations of the GSF with high accuracy will require a huge amount of time and computer resources mainly because of the regularization problem induced by the point mass limit. Therefore it is important to develop a way to reduce the cost of computing the GSF. The two-timescale expansion method [6] gives a hint for it: assuming that a point mass does not encounter any transient resonances (*e.g.* shown in [7]), the orbital phase, which is the most important information to predict the waveform, can be expressed in the expansion with respect to the mass ratio, η , as

$$\Phi = \eta^{-1} \left[\Phi^{(0)} + \eta \Phi^{(1)} + O(\eta^2) \right], \quad (1)$$

where $\Phi^{(0)}$ and $\Phi^{(1)}$ are quantities of order unity. The leading term, $\Phi^{(0)}$, can be calculated from the knowledge up to the time-averaged dissipative piece of the first order GSF, corresponding to the secular growth. The calculation of this secular contribution can be simplified significantly by using the radiative field defined as half the retarded solution minus half the advanced solution for the equation of the gravitational perturbation [8–10], *i.e.* the adiabatic approximation method, because the radiative field is the homogeneous solution free from the divergence induced by the point mass limit. This method allows us to calculate the leading term accurately without spending huge computational resources. On the other hand, the calculation of $\Phi^{(1)}$ requires the rest of the first order GSF (the oscillatory part of the dissipative GSF and the conservative GSF) and the time-averaged dissipative piece of the second order GSF. There is no simplification in calculating these post-1 adiabatic pieces at present. Since $\Phi^{(1)}$ is subleading, however, the requirement of the accuracy is not so high compared to that of the leading term. This fact suggests that it is possible to reduce the computational cost by using a suitable method with an appropriate error tolerance to calculate each piece of the GSF (for example, a hybrid approach is proposed in [11]).

In this work, we focus on the time-averaged dissipative part of the first order GSF, which has the dominant contribution to the evolution of inspirals, and present the analytic post-Newtonian (PN) formulae. So far, several works in this direction had been done for two restricted classes of orbits: circular orbits and equatorial orbits. (See [12] and references therein for early works in 1990's). Recently, thanks to the progress of computer technology, the very higher order post-Newtonian calculations can be possible for circular equatorial orbits: the 22PN calculation of the energy flux is demonstrated in Schwarzschild case [13], and the 11PN calculation in Kerr case [14]. There is also the calculation of the secular GSF effects for slightly eccentric and slightly inclined (non-equatorial) orbits [10], and later it had been extended to orbits with arbitrary inclination [15], where the PN formulae of the secular GSF effects are presented in the expansion with respect to the orbital eccentricity. However, the calculation in [15] has been done only up to the 2.5PN order with the second order correction of the eccentricity. Also the absorption to the black hole is ignored there. The main purpose of this work is to update the results in [15] up to the 4PN order and the sixth order correction of the eccentricity, including the effect of the absorption to the black hole.

This paper is organized as follows. In Sec.2, we give a brief review of the geodesic motion of a point particle in Kerr spacetime, the gravitational perturbations induced by the particle, and the adiabatic approximation method of calculating the secular effect of the GSF. In Sec.3.1, we present the PN formulae of the secular changes of the energy, azimuthal angular momentum, and Carter parameter of the particle due to the gravitational radiation reaction in the expansion with respect to the orbital eccentricity. In Sec.3.2, we investigate the accuracy of our PN formulae by comparing to numerical results given by the method in [16–18], which can give each modal flux at the accuracy about 14 significant figures. In Sec.3.3, we implement a resummation method to the PN formulae given in this work in order to improve the accuracy. In Sec.3.4, we discuss the convergence of the analytic formulae as the PN expansion and the expansion with respect to the eccentricity. Finally, we summarize the paper in Sec.4. For the readability of the main text, we present the PN formulae for the orbital parameters, the fundamental frequencies, the orbital motion in Appendices A and B, which are used in calculating the secular changes of the orbital parameters. We also present the PN formulae for the secular changes of an alternative set of the orbital parameters in Appendix C. Throughout this paper we use metric signature $(-+++)$ and “geometrized” units with $c = G = 1$.

2. Review of formulation: Adiabatic radiation reaction

The orbital evolution of a point particle due to the time-averaged dissipative part of the GSF is often described in terms of the secular changes of the orbital parameters. In order to calculate the changes, we need the information on the first order gravitational perturbations induced by the particle when it moves along the background geodesics. In this section, we review the geodesic dynamics of a point particle in Kerr spacetime, the gravitational perturbations induced by the particle, and the adiabatic evolution of the orbital parameters.

2.1. Geodesic motion

The Kerr metric in the Boyer-Lindquist coordinates, (t, r, θ, φ) , is given by

$$g_{\mu\nu}dx^\mu dx^\nu = -\left(1 - \frac{2Mr}{\Sigma}\right)dt^2 - \frac{4Mar\sin^2\theta}{\Sigma}dtd\varphi + \frac{\Sigma}{\Delta}dr^2 + \Sigma d\theta^2 + \left(r^2 + a^2 + \frac{2Ma^2r}{\Sigma}\sin^2\theta\right)\sin^2\theta d\varphi^2, \quad (2)$$

where $\Sigma = r^2 + a^2\cos^2\theta$, $\Delta = r^2 - 2Mr + a^2$, M and aM are the mass and angular momentum of the black hole, respectively.

There are two Killing vectors related to the stationarity and axisymmetry of Kerr spacetime, which are expressed as $\xi_{(t)}^\mu = (1, 0, 0, 0)$ and $\xi_{(\varphi)}^\mu = (0, 0, 0, 1)$. In addition, it is known that Kerr spacetime possesses a Killing tensor, $K_{\mu\nu} = 2\Sigma l_{(\mu}n_{\nu)} + r^2g_{\mu\nu}$, where l^μ and n^μ are the Kinnersley’s null vectors given by

$$l^\mu := \left(\frac{r^2 + a^2}{\Delta}, 1, 0, \frac{a}{\Delta}\right), \quad n^\mu := \left(\frac{r^2 + a^2}{2\Sigma}, -\frac{\Delta}{2\Sigma}, 0, \frac{a}{2\Sigma}\right). \quad (3)$$

For the geodesic motion of a particle in Kerr geometry, there are three constants of motion related to the symmetries:

$$\hat{E} := -u^\alpha \xi_\alpha^{(t)}, \quad \hat{L} := u^\alpha \xi_\alpha^{(\varphi)}, \quad \hat{Q} := K_{\alpha\beta}u^\alpha u^\beta, \quad (4)$$

where u^α is the four velocity of the particle. $\hat{E}$ and $\hat{L}$ correspond to the specific energy and azimuthal angular momentum of the particle respectively. $\hat{Q}$ is called as the Carter constant, which corresponds to the square of the specific total angular momentum in Schwarzschild case. These specific variables are measured in units of the particle's mass, μ . One can recover the expressions in the standard units as

$$E := \mu \hat{E}, \quad L := \mu \hat{L}, \quad Q := \mu^2 \hat{Q}. \quad (5)$$

There is another definition of the Carter constant, $C \equiv Q - (aE - L)^2$, which vanishes for equatorial orbits. In this paper, we use C as one of the orbital parameters, instead of Q .

By using these constants of motion, the geodesic equations can be expressed in the following form as

$$\left(\frac{dr}{d\lambda}\right)^2 = R(r), \quad \left(\frac{d\cos\theta}{d\lambda}\right)^2 = \Theta(\cos\theta), \quad (6)$$

$$\frac{dt}{d\lambda} = V_{tr}(r) + V_{t\theta}(\theta), \quad \frac{d\varphi}{d\lambda} = V_{\varphi r}(r) + V_{\varphi\theta}(\theta), \quad (7)$$

where we introduced a new parameter λ through the relation $d\lambda = d\tau/\Sigma$, and some functions as

$$P(r) := \hat{E}(r^2 + a^2) - a\hat{L}, \quad (8)$$

$$R(r) := [P(r)]^2 - \Delta[r^2 + (a\hat{E} - \hat{L})^2 + \hat{C}], \quad (9)$$

$$\Theta(\cos\theta) := \hat{C} - (\hat{C} + a^2(1 - \hat{E}^2) + \hat{L}^2)\cos^2\theta + a^2(1 - \hat{E}^2)\cos^4\theta, \quad (10)$$

$$V_{tr}(r) := \frac{r^2 + a^2}{\Delta}P(r), \quad V_{t\theta}(\theta) := -a(a\hat{E}\sin^2\theta - \hat{L}), \quad (11)$$

$$V_{\varphi r}(r) := \frac{a}{\Delta}P(r), \quad V_{\varphi\theta}(\theta) := -\left(a\hat{E} - \frac{\hat{L}}{\sin^2\theta}\right). \quad (12)$$

A generic geodesic orbit in Kerr spacetime can be characterized by three parameters, $\{E, L, C\}$ ¹. In the case of a bound orbit, we can use an alternative set of parameters, $\{r_p, r_a, \theta_{\min}\}$, instead of $\{E, L, C\}$, where r_p and r_a are the values of r at the periapsis and apoapsis and $\theta_{\min}$ is the minimal value of θ , respectively. Using this set of parameters, we can describe the range in which the motion takes place as $r_p \leq r \leq r_a$ and $\theta_{\min} \leq \theta \leq \pi - \theta_{\min}$. There is another useful choice of parameters used in [19], $\{p, e, \iota\}$, defined by

$$p := \frac{2r_p r_a}{M(r_a + r_p)}, \quad e := \frac{r_a - r_p}{r_a + r_p}, \quad \cos\iota := \frac{L}{\sqrt{L^2 + C}}. \quad (13)$$

By analogy to the parametrization used in celestial mechanics, p , e , ι are referred as semi-latus rectum, orbital eccentricity, orbital inclination angle, respectively. For later convenience, we also introduce $Y = \cos\iota$ and $v = \sqrt{1/p}$. Since v corresponds to the magnitude of the orbital velocity, it can be used as the post-Newtonian parameter. For example, we

¹ Strictly speaking, the orbit is also characterized by the initial position of the particle. However, the time-averaged dissipative part of the first order GSF does not affect the initial position (the other parts of the first order GSF and the higher order GSF will do) [6]. Also the secular changes of $\{E, L, C\}$ does not depend on the initial position. Hence we do not need the information on the initial position to describe the secular evolution of the orbit at the order considered in this paper.

call the $O(v^8)$ -correction from the leading order as the fourth order post-Newtonian (4PN) correction.

It is worth noting that, by introducing λ , the radial and longitudinal equations of motion in Eq.(6), are completely decoupled. For an bound orbit, therefore, the radial and longitudinal motions are periodic with the periods, $\{\Lambda_r, \Lambda_\theta\}$, defined by

$$\Lambda_r = 2 \int_{r_p}^{r_a} \frac{dr}{\sqrt{R(r)}}, \quad \Lambda_\theta = 4 \int_{\theta_{\min}}^{\pi/2} \frac{d\theta}{\sqrt{\Theta(\theta)}}. \quad (14)$$

This means that these motions can be expressed in terms of Fourier series as

$$r(\lambda) = p \sum_{n_r=0}^{\infty} \alpha_{n_r} \cos n_r \Omega_r \lambda, \quad (15)$$

$$\cos \theta(\lambda) = \sqrt{1 - Y^2} \sum_{n_\theta=0}^{\infty} \beta_{n_\theta} \sin n_\theta \Omega_\theta \lambda, \quad (16)$$

where Ω_r and Ω_θ are the radial and longitudinal frequencies given by

$$\Omega_r := \frac{2\pi}{\Lambda_r}, \quad \Omega_\theta := \frac{2\pi}{\Lambda_\theta}, \quad (17)$$

and we choose the initial values so that $r(\lambda = 0) = r_p$ and $\theta(\lambda = 0) = \pi/2$.²

Since the temporal and azimuthal equations of motion in Eq.(7) are divided into the r - and θ -dependent parts, the solutions can be divided into three parts: the linear term with respect to λ , the oscillatory part with period of Λ_r , and the oscillatory part with period of Λ_θ . They can be expressed as

$$t(\lambda) = \Omega_t \lambda + t^{(r)}(\lambda) + t^{(\theta)}(\lambda); \quad t^{(A)}(\lambda) := \sum_{n_A=1}^{\infty} \tilde{t}_{n_A}^{(A)} \sin n_A \Omega_A \lambda, \quad (18)$$

$$\varphi(\lambda) = \Omega_\varphi \lambda + \varphi^{(r)}(\lambda) + \varphi^{(\theta)}(\lambda); \quad \varphi^{(A)}(\lambda) := \sum_{n_A=1}^{\infty} \tilde{\varphi}_{n_A}^{(A)} \sin n_A \Omega_A \lambda, \quad (19)$$

where the index A runs over $\{r, \theta\}$, and

$$\Omega_t := \left\langle \frac{dt}{d\lambda} \right\rangle_\lambda, \quad \Omega_\varphi := \left\langle \frac{d\varphi}{d\lambda} \right\rangle_\lambda \quad (20)$$

with $\langle \cdots \rangle_\lambda \equiv \lim_{T \rightarrow \infty} (2T)^{-1} \int_{-T}^T d\lambda \cdots$, representing the time average along the geodesic. We choose the initial conditions as $t(\lambda = 0) = \varphi(\lambda = 0) = 0$. Ω_φ corresponds to the frequency of the orbital rotation.

In Appendices A and B, we present the PN formulae of the orbital parameters, $\{E, L, C\}$, the fundamental frequencies, $\{\Omega_t, \Omega_r, \Omega_\theta, \Omega_\phi\}$, and the Fourier coefficients of the motions in Eqs. (15), (16), (18) and (19).

² If the ratio of the radial and longitudinal frequencies is irrational, we can adjust the origin of λ approximately so that the radial and longitudinal oscillations reach the minima simultaneously at $\lambda = 0$ [8]. On the other hand, it is not the case if the ratio is rational, *i.e.* the resonance case. This implies that the secular evolution of a resonant orbit cannot be described only by the PN formulae derived in this work [20].

2.2. Secular evolution of orbital parameters

The gravitational perturbations in Kerr spacetime can be described by the Weyl scalar, Ψ_4 , which satisfies the Teukolsky equation [21]. To solve the Teukolsky equation, the method of separation of variables is often used, in which Ψ_4 is decomposed in the form as

$$\Psi_4 = \sum_{\ell m} \int d\omega R_\Lambda(r) S_\Lambda(\theta) e^{im\varphi - i\omega t}, \quad (21)$$

where $S_\Lambda(\theta)$ is the spin-2 spheroidal harmonics and Λ represents a set of indices in the Fourier-harmonic expansion, $\{\ell, m, \omega\}$. The separated equation for the radial function is given by

$$\left[\Delta^2 \frac{d}{dr} \left(\Delta^{-1} \frac{d}{dr} \right) + \left(\frac{K^2 + 4i(r-M)K}{\Delta} - 8i\omega r - \bar{\lambda} \right) \right] R_\Lambda(r) = T_\Lambda, \quad (22)$$

where

$$K \equiv (r^2 + a^2)\omega - ma,$$

T_Λ is the source term constructed from the energy-momentum tensor of the point particle, and $\bar{\lambda}$ is the eigenvalue determined by the equation for S_Λ (To find the basic formulae for the Teukolsky formalism used in this paper, refer to the section 2 in [22] for example).

The amplitudes of the partial waves at the horizon and at infinity are defined by the asymptotic forms of the solution of the radial equation as

$$R_\Lambda(r \rightarrow r_+) \equiv \mu Z_\Lambda^H \Delta^2 e^{-ikr^*}, \quad R_\Lambda(r \rightarrow \infty) \equiv \mu Z_\Lambda^\infty r^3 e^{i\omega r^*} \quad (23)$$

with $r_+ \equiv M + \sqrt{M^2 - a^2}$ and $k = \omega - ma/(2Mr_+)$. Since the spectrum with respect to ω gets discrete in the case of a bound orbit, $Z_\Lambda^{H,\infty}$ take the form

$$Z_\Lambda^{H,\infty} = 2\pi\delta(\omega - \omega_{mn_r n_\theta}) \tilde{Z}_\Lambda^{H,\infty}, \quad (24)$$

where $\tilde{\Lambda}$ denotes the set of indices, $\{\ell, m, n_r, n_\theta\}$, and

$$\omega_{mn_r n_\theta} \equiv \Omega_t^{-1} (m\Omega_\varphi + n_r\Omega_r + n_\theta\Omega_\theta). \quad (25)$$

With these amplitudes, the secular changes of the orbital parameters, $\{E, L, C\}$, can be expressed by

$$\left\langle \frac{dE}{dt} \right\rangle_t = -\mu^2 \sum_{\tilde{\Lambda}} \frac{1}{4\pi\omega_{mn_r n_\theta}^2} \left(\left| \tilde{Z}_\Lambda^\infty \right|^2 + \alpha_{\ell m}(\omega_{mn_r n_\theta}) \left| \tilde{Z}_\Lambda^H \right|^2 \right), \quad (26)$$

$$\left\langle \frac{dL}{dt} \right\rangle_t = -\mu^2 \sum_{\tilde{\Lambda}} \frac{m}{4\pi\omega_{mn_r n_\theta}^3} \left(\left| \tilde{Z}_\Lambda^\infty \right|^2 + \alpha_{\ell m}(\omega_{mn_r n_\theta}) \left| \tilde{Z}_\Lambda^H \right|^2 \right), \quad (27)$$

$$\begin{aligned} \left\langle \frac{dC}{dt} \right\rangle_t &= -2 \langle a^2 E \cos^2 \theta \rangle_\lambda \left\langle \frac{dE}{dt} \right\rangle_t + 2 \langle L \cot^2 \theta \rangle_\lambda \left\langle \frac{dL}{dt} \right\rangle_t \\ &\quad - \mu^3 \sum_{\tilde{\Lambda}} \frac{n_\theta \Omega_\theta}{2\pi\omega_{mn_r n_\theta}^3} \left(\left| \tilde{Z}_\Lambda^\infty \right|^2 + \alpha_{\ell m}(\omega_{mn_r n_\theta}) \left| \tilde{Z}_\Lambda^H \right|^2 \right), \end{aligned} \quad (28)$$

where

$$\alpha_{\ell m}(\omega) = \frac{256(2Mr_+)^5 k(k^2 + 4\tilde{\epsilon}^2)(k^2 + 16\tilde{\epsilon}^2)\omega^3}{|\mathcal{C}_S|^2}, \quad \tilde{\epsilon} = \sqrt{M^2 - a^2}/(4Mr_+), \quad (29)$$

and $\mathcal{C}_S$ is the Starobinsky constant given by [23]

$$|\mathcal{C}_S|^2 = [(\bar{\lambda} + 2)^2 + 4a\omega m - 4a^2\omega^2] [\bar{\lambda}^2 + 36a\omega m - 36a^2\omega^2]$$

$$+(2\bar{\lambda} + 3)(96a^2\omega^2 - 48a\omega m) + 144\omega^2(M^2 - a^2). \quad (30)$$

It should be noted that, in these formulae, the averaged rates of change are expressed with respect to the Boyer-Lindquist time, which can be related to those with respect to λ [24] as

$$\left\langle \frac{dI}{dt} \right\rangle_t = \left\langle \frac{dt}{d\lambda} \right\rangle_\lambda^{-1} \left\langle \frac{dI}{d\lambda} \right\rangle_\lambda \quad (31)$$

for a function of time, $I(t)$. Also it should be noted that each formula in Eqs. (26)-(28) can be divided into the infinity part and the horizon part: the former consists of the terms including the amplitudes of the partial waves at the infinity, $\tilde{Z}_\Lambda^\infty$, the latter consists of the terms including the amplitudes at the horizon, $\tilde{Z}_\Lambda^H$. As for the energy and azimuthal angular momentum, the infinity parts are balanced with the corresponding fluxes radiated to infinity and the horizon parts with the absorption of the gravitational waves into the central black hole [23, 25].

The practical calculation of $\tilde{Z}_\Lambda^{H,\infty}$ involves solving the geodesic equations, calculating two independent homogeneous solutions of Eq.(22) and the spin-2 spheroidal harmonics, and the Fourier transformation of functions consisting of them. In this work, we followed the same procedure proposed in [15] to perform these calculations analytically.

In performing the summation in Eqs. (26)-(28) practically, we need to truncate the summation to finite ranges of $\tilde{\Lambda} = \{\ell, m, n_r, n_\theta\}$. To obtain the accuracy of the 4PN and $O(e^6)$, it is necessary to sum ℓ in the range $2 \leq \ell \leq 6$ ($2 \leq \ell \leq 3$), n_r in the range $-3 \leq n_r \leq 3$ ($-2 \leq n_r \leq 3$) and n_θ in the range $-8 \leq n_\theta \leq 12$ ($-4 \leq n_\theta \leq 6$) for the infinity (horizon) part. The other modes out of these ranges are the higher PN corrections than the 4PN order or the higher order corrections than $O(e^6)$.

3. Results

3.1. PN formulae of the secular changes of orbital parameters

In this work, we derived the analytic 4PN order formulae of Eqs.(26)-(28) in the expansion with respect to the orbital eccentricity, e , up to $O(e^6)$ (we simply call them as the 4PN $O(e^6)$ formulae). Since the full expressions of the 4PN $O(e^6)$ formulae are too lengthy to show in the text, we show the infinity parts up to the 3PN order and the horizon parts up to the 3.5PN order (while we keep the expansions with respect to e up to $O(e^6)$). The complete expressions of the 4PN $O(e^6)$ formulae will be publicly available online [26].

The infinity parts of Eqs.(26)-(28) are given by

$$\begin{aligned} \left\langle \frac{dE}{dt} \right\rangle_t^\infty &= \left(\frac{dE}{dt} \right)_N \left[1 + \frac{73}{24}e^2 + \frac{37}{96}e^4 + \left\{ -\frac{1247}{336} - \frac{9181}{672}e^2 + \frac{809}{128}e^4 + \frac{8609}{5376}e^6 \right\} v^2 \right. \\ &\quad + \left\{ 4\pi - \frac{73}{12}Yq + \left(\frac{1375}{48}\pi - \frac{823}{24}Yq \right) e^2 \right. \\ &\quad \left. + \left(\frac{3935}{192}\pi - \frac{949}{32}Yq \right) e^4 + \left(\frac{10007}{9216}\pi - \frac{491}{192}Yq \right) e^6 \right\} v^3 \\ &\quad + \left\{ -\frac{44711}{9072} + \frac{527}{96}Y^2q^2 - \frac{329}{96}q^2 + \left(-\frac{172157}{2592} - \frac{4379}{192}q^2 + \frac{6533}{192}Y^2q^2 \right) e^2 \right. \\ &\quad \left. + \left(-\frac{2764345}{24192} - \frac{3823}{256}q^2 + \frac{6753}{256}Y^2q^2 \right) e^4 \right\} v^4 \end{aligned}$$

$$\begin{aligned}
& + \left(\frac{3743}{2304} - \frac{363}{512} q^2 + \frac{2855}{1536} Y^2 q^2 \right) e^6 \Big\} v^4 \\
& + \left\{ \frac{3749}{336} Y q - \frac{8191}{672} \pi + \left(-\frac{44531}{336} \pi + \frac{1759}{56} Y q \right) e^2 \right. \\
& \quad \left. - \left(\frac{4311389}{43008} \pi + \frac{111203}{1344} Y q \right) e^4 + \left(\frac{15670391}{387072} \pi - \frac{49685}{448} Y q \right) e^6 \right\} v^5 \\
& + \left\{ \frac{6643739519}{69854400} - \frac{1712}{105} \gamma - \frac{3424}{105} \ln(2) + \frac{16}{3} \pi^2 + \frac{135}{8} q^2 - \frac{169}{6} \pi Y q \right. \\
& \quad + \frac{73}{21} Y^2 q^2 + \left(\frac{43072561991}{27941760} + \frac{680}{9} \pi^2 - \frac{234009}{560} \ln(3) - \frac{14552}{63} \gamma \right. \\
& \quad \left. \left. - \frac{13696}{315} \ln(2) + \frac{205747}{1344} q^2 - \frac{4339}{16} \pi Y q + \frac{13697}{192} Y^2 q^2 \right) e^2 \right. \\
& \quad + \left(\frac{919773569303}{279417600} + \frac{5171}{36} \pi^2 + \frac{2106081}{448} \ln(3) - \frac{12295049}{1260} \ln(2) \right. \\
& \quad \left. - \frac{553297}{1260} \gamma + \frac{208571}{1792} q^2 - \frac{42271}{96} \pi Y q + \frac{471711}{1792} Y^2 q^2 \right) e^4 \\
& \quad + \left(\frac{308822406727}{186278400} - \frac{864819261}{35840} \ln(3) - \frac{187357}{1260} \gamma + \frac{1751}{36} \pi^2 \right. \\
& \quad \left. - \frac{5224609375}{193536} \ln(5) + \frac{24908851}{252} \ln(2) + \frac{3253}{10752} q^2 - \frac{4867907}{27648} \pi Y q \right. \\
& \quad \left. + \frac{289063}{1536} Y^2 q^2 \right) e^6 \\
& \quad \left. - \left(\frac{1712}{105} + \frac{14552}{63} e^2 + \frac{553297}{1260} e^4 + \frac{187357}{1260} e^6 \right) \ln v \right\} v^6 \Big], \tag{32}
\end{aligned}$$

$$\begin{aligned}
\left\langle \frac{dL}{dt} \right\rangle_t^\infty &= \left(\frac{dL}{dt} \right)_N \left[\left\{ 1 + \frac{7}{8} e^2 \right\} Y + \left\{ -\frac{1247}{336} - \frac{425}{336} e^2 + \frac{10751}{2688} e^4 \right\} Y v^2 \right. \\
& + \left\{ \frac{61}{24} q - \frac{61}{8} Y^2 q + 4 \pi Y + \left(\frac{63}{8} q + \frac{97}{8} \pi Y - \frac{91}{4} Y^2 q \right) e^2 \right. \\
& \quad \left. + \left(\frac{95}{64} q + \frac{49}{32} \pi Y - \frac{461}{64} Y^2 q \right) e^4 - \frac{49}{4608} \pi Y e^6 \right\} v^3 \\
& + \left\{ -\frac{44711}{9072} - \frac{57}{16} q^2 + \frac{45}{8} Y^2 q^2 + \left(-\frac{302893}{6048} - \frac{201}{16} q^2 + \frac{37}{2} Y^2 q^2 \right) e^2 \right. \\
& \quad \left. + \left(-\frac{701675}{24192} - \frac{351}{128} q^2 + \frac{331}{64} Y^2 q^2 \right) e^4 + \frac{162661}{16128} e^6 \right\} Y v^4 \\
& + \left\{ \frac{4301}{224} Y^2 q - \frac{8191}{672} \pi Y - \frac{2633}{224} q \right. \\
& \quad + \left(-\frac{66139}{1344} q - \frac{48361}{1344} \pi Y + \frac{18419}{448} Y^2 q \right) e^2 \\
& \quad + \left(\frac{3959}{1792} q + \frac{1657493}{43008} \pi Y - \frac{257605}{5376} Y^2 q \right) e^4 \\
& \quad \left. + \left(\frac{19161}{3584} q + \frac{5458969}{774144} \pi Y - \frac{52099}{1536} Y^2 q \right) e^6 \right\} v^5
\end{aligned}$$

$$\begin{aligned}
& + \left\{ \frac{145}{12} \pi q + \frac{6643739519}{69854400} Y + \frac{16}{3} \pi^2 Y - \frac{1712}{105} \gamma Y - \frac{3424}{105} \ln(2) Y \right. \\
& \quad - \frac{171}{112} Y q^2 - \frac{145}{4} \pi Y^2 q + \frac{1769}{112} Y^3 q^2 \\
& \quad + \left(\frac{995}{12} \pi q + \frac{229}{6} \pi^2 Y + \frac{6769212511}{8731800} Y + \frac{1391}{30} \ln(2) Y - \frac{24503}{210} \gamma Y \right. \\
& \quad \quad \left. - \frac{78003}{280} \ln(3) Y - \frac{46867}{1344} Y q^2 - \frac{877}{4} \pi Y^2 q + \frac{27997}{192} Y^3 q^2 \right) e^2 \\
& \quad + \left(\frac{21947}{384} \pi q + \frac{4795392143}{7761600} Y + \frac{3042117}{1120} \ln(3) Y - \frac{418049}{84} \ln(2) Y \right. \\
& \quad \quad - \frac{11663}{140} \gamma Y + \frac{109}{4} \pi^2 Y - \frac{1481}{16} Y q^2 - \frac{22403}{128} \pi Y^2 q + \frac{267563}{1344} Y^3 q^2 \left. \right) e^4 \\
& \quad + \left(\frac{38747}{13824} \pi q + \frac{31707715321}{186278400} Y + \frac{23}{16} \pi^2 Y + \frac{94138279}{2160} \ln(2) Y \right. \\
& \quad \quad - \frac{1044921875}{96768} \ln(5) Y - \frac{42667641}{3584} \ln(3) Y - \frac{2461}{560} \gamma Y - \frac{68333}{3584} Y q^2 \\
& \quad \quad \left. - \frac{59507}{4608} \pi Y^2 q + \frac{183909}{3584} Y^3 q^2 \right) e^6 \\
& \quad \left. - \left(\frac{1712}{105} + \frac{24503}{210} e^2 + \frac{11663}{140} e^4 + \frac{2461}{560} e^6 \right) Y \ln v \right\} v^6 \Big], \tag{33}
\end{aligned}$$

$$\begin{aligned}
\left\langle \frac{dC}{dt} \right\rangle_t^\infty &= \left(\frac{dC}{dt} \right)_N \left[1 + \frac{7}{8} e^2 + \left(-\frac{743}{336} + \frac{23}{42} e^2 + \frac{11927}{2688} e^4 \right) v^2 \right. \\
& + \left\{ 4\pi - \frac{85}{8} Y q + \left(\frac{97}{8} \pi - \frac{211}{8} Y q \right) e^2 \right. \\
& \quad \left. + \left(\frac{49}{32} \pi - \frac{517}{64} Y q \right) e^4 - \frac{49}{4608} \pi e^6 \right\} v^3 \\
& + \left\{ -\frac{129193}{18144} - \frac{329}{96} q^2 + \frac{53}{8} Y^2 q^2 + \left(-\frac{84035}{1728} - \frac{929}{96} q^2 + \frac{163}{8} Y^2 q^2 \right) e^2 \right. \\
& \quad \left. + \left(-\frac{1030273}{48384} - \frac{1051}{768} q^2 + \frac{387}{64} Y^2 q^2 \right) e^4 + \frac{100103}{8064} e^6 \right\} v^4 \\
& + \left\{ -\frac{4159}{672} \pi + \frac{2553}{224} Y q + \left(-\frac{21229}{1344} \pi - \frac{553}{192} Y q \right) e^2 \right. \\
& \quad \left. + \left(\frac{2017013}{43008} \pi - \frac{475541}{5376} Y q \right) e^4 + \left(\frac{6039325}{774144} \pi - \frac{153511}{3584} Y q \right) e^6 \right\} v^5 \\
& + \left\{ \frac{11683501663}{139708800} + \frac{16}{3} \pi^2 - \frac{1712}{105} \gamma - \frac{3424}{105} \ln(2) + \frac{1277}{192} q^2 - \frac{193}{4} \pi Y q \right. \\
& \quad + \frac{2515}{48} Y^2 q^2 + \left(\frac{16319179321}{23284800} + \frac{229}{6} \pi^2 - \frac{24503}{210} \gamma + \frac{1391}{30} \ln(2) \right. \\
& \quad \quad \left. - \frac{78003}{280} \ln(3) + \frac{16979}{1344} q^2 - \frac{2077}{8} \pi Y q + \frac{118341}{448} Y^2 q^2 \right) e^2 \\
& \quad \left. + \left(\frac{211889615389}{372556800} + \frac{109}{4} \pi^2 + \frac{3042117}{1120} \ln(3) - \frac{11663}{140} \gamma - \frac{418049}{84} \ln(2) \right. \right.
\end{aligned}$$

$$\begin{aligned}
& -\frac{132193}{3584} q^2 - \frac{24543}{128} \pi Y q + \frac{91747}{336} Y^2 q^2 \Big) e^4 \\
& + \left(\frac{33928992071}{186278400} - \frac{1044921875}{96768} \ln(5) + \frac{23}{16} \pi^2 - \frac{42667641}{3584} \ln(3) \right. \\
& \quad \left. + \frac{94138279}{2160} \ln(2) - \frac{2461}{560} \gamma - \frac{24505}{5376} q^2 - \frac{4151}{288} \pi Y q + \frac{718799}{10752} Y^2 q^2 \right) e^6 \\
& - \left(\frac{1712}{105} + \frac{24503}{210} e^2 + \frac{11663}{140} e^4 + \frac{2461}{560} e^6 \right) \ln v \Big\} v^6 \Big], \tag{34}
\end{aligned}$$

where the leading contributions are given by

$$\begin{aligned}
\left(\frac{dE}{dt} \right)_N &= -\frac{32}{5} \left(\frac{\mu}{M} \right)^2 v^{10} (1 - e^2)^{3/2}, \\
\left(\frac{dL}{dt} \right)_N &= -\frac{32}{5} \left(\frac{\mu^2}{M} \right) v^7 (1 - e^2)^{3/2}, \\
\left(\frac{dC}{dt} \right)_N &= -\frac{64}{5} \mu^3 v^6 (1 - e^2)^{3/2} (1 - Y^2). \tag{35}
\end{aligned}$$

The horizon parts of Eqs.(26)-(28) are given by

$$\begin{aligned}
\left\langle \frac{dE}{dt} \right\rangle_t^H &= \left(\frac{dE}{dt} \right)_N \left[-\frac{1}{512} (16 + 120 e^2 + 90 e^4 + 5 e^6) (8 + 9 q^2 + 15 Y^2 q^2) q Y v^5 \right. \\
& - \left\{ 1 + \frac{81}{32} q^2 - \frac{15}{32} Y^2 q^2 + \left(\frac{57}{4} + \frac{1143}{32} q^2 - \frac{195}{32} Y^2 q^2 \right) e^2 \right. \\
& \quad + \left(\frac{465}{16} + \frac{4455}{64} q^2 - \frac{225}{32} Y^2 q^2 \right) e^4 \\
& \quad \left. \left. + \left(\frac{355}{32} + \frac{6345}{256} q^2 + \frac{75}{256} Y^2 q^2 \right) e^6 \right\} q Y v^7 \right], \tag{36}
\end{aligned}$$

$$\begin{aligned}
\left\langle \frac{dL}{dt} \right\rangle_t^H &= \left(\frac{dL}{dt} \right)_N \left[-\frac{(8 + 24 e^2 + 3 e^4)}{1024} (16 + 33 q^2 + 16 Y^2 + 18 Y^2 q^2 + 45 Y^4 q^2) q v^5 \right. \\
& - \left\{ \frac{5}{4} + \frac{375}{128} q^2 - \frac{1}{4} Y^2 - \frac{63}{64} Y^2 q^2 + \frac{15}{128} Y^4 q^2 \right. \\
& \quad + \left(10 + \frac{5955}{256} q^2 - \frac{5}{4} Y^2 - \frac{855}{128} Y^2 q^2 + \frac{675}{256} Y^4 q^2 \right) e^2 \\
& \quad + \left(\frac{255}{32} + \frac{18855}{1024} q^2 - \frac{15}{32} Y^2 - \frac{2295}{512} Y^2 q^2 + \frac{3375}{1024} Y^4 q^2 \right) e^4 \\
& \quad \left. \left. + \left(\frac{15}{32} + \frac{2205}{2048} q^2 - \frac{225}{1024} Y^2 q^2 + \frac{525}{2048} Y^4 q^2 \right) e^6 \right\} q v^7 \right], \tag{37}
\end{aligned}$$

$$\begin{aligned}
\left\langle \frac{dC}{dt} \right\rangle_t^H &= \left(\frac{dC}{dt} \right)_N \left[-\frac{1}{1024} (8 + 24 e^2 + 3 e^4) (16 + 3 q^2 + 45 Y^2 q^2) q Y v^5 \right. \\
& + \left\{ \frac{1}{16} + \frac{93}{256} q^2 - \frac{165}{256} Y^2 q^2 + \left(\frac{5}{8} + \frac{705}{256} q^2 - \frac{1125}{256} Y^2 q^2 \right) e^2 \right. \\
& \quad \left. + \left(\frac{27}{128} + \frac{4131}{2048} q^2 - \frac{8235}{2048} Y^2 q^2 \right) e^4 \right. \\
& \quad \left. \left. + \left(\frac{27}{128} + \frac{4131}{2048} q^2 - \frac{8235}{2048} Y^2 q^2 \right) e^4 \right\} q Y v^7 \right],
\end{aligned}$$

$$+ \left(-\frac{3}{128} + \frac{27}{256} q^2 - \frac{165}{512} Y^2 q^2 \right) e^6 \Big\} q Y v^7 \Big]. \quad (38)$$

$\langle dE/dt \rangle_t^H$, $\langle dL/dt \rangle_t^H$ and $\langle dC/dt \rangle_t^H$ in Eqs. (36)-(38) are new PN formulae derived in this paper. $\langle dE/dt \rangle_t^\infty$, $\langle dL/dt \rangle_t^\infty$ and $\langle dC/dt \rangle_t^\infty$ in Eqs. (32)-(34) are consistent with those in Ref. [15] up to 2.5PN and $O(e^2)$.

From the leading order expressions in Eq. (35), one will find that the Carter parameter, C , does not change due to the radiation of the gravitational waves when $Y = 1$ (equatorial orbits) because $(dC/dt)_N = 0$.

In the Schwarzschild case, the Carter parameter corresponds to the square of the equatorial angular momentum (the normal component to the rotational axis of the central black hole). Then there is expected to exist the duality between L^2 and C due to the spherical symmetry. In fact, from Eqs. (33) and (34), (and also from (37) and (38)), one can find that $\langle dL^2/dt \rangle_t$ vanishes in $Y = 0$ (polar orbits) while $\langle dC/dt \rangle_t$ for $Y = 0$ coincides with $\langle dL^2/dt \rangle_t$ for $Y = 1$. This can be also realized by seeing that the secular change of the total angular momentum, $\langle d(L^2 + C)/dt \rangle_t$, is independent of Y . Then, it might be possible to understand that $\langle dL/dt \rangle_t^\infty$ becomes 1.5PN from the leading order when $q \neq 0$ and $Y = 0$ (polar orbits) due to the spin-orbit coupling.

From the expressions of the horizon parts shown in Eqs. (36)-(38), we find that the absorption of the gravitational waves to the central black hole contributes at $O(v^5)$ from the leading order in Eq. (35) for $q \neq 0$ and at $O(v^8)$ for $q = 0$. The $O(v^5)$ and $O(v^7)$ corrections in $\langle dE/dt \rangle_t^H$ can be positive for $q > 0$, which means that the particle can gain the energy through a superradiance phenomenon. These observations are consistent with the results for circular, equatorial orbits shown in Refs. [27, 28].

We also find that the superradiance terms in Eq. (36) vanish for $Y = 0$, and that $\langle dE/dt \rangle_t^H$ has only the 4PN and higher order corrections. The superradiance terms may come from the coupling between the black hole spin and the orbital angular momentum, like $\propto \mathbf{L} \cdot \mathbf{S} \propto q \cos \iota$. Hence, when the orbital inclination increases (Y gets small correspondingly), the superradiance is suppressed [29].

3.2. Comparison to numerical results

To investigate the accuracy of the 4PN $O(e^6)$ formulae derived in this work, we compare them to the corresponding numerical results given by the method established in Ref. [16–18], which enables one to compute the modal fluxes with the relative error of $\sim 10^{-14}$ in double precision computations. In the practical computations, as well as in deriving the analytic expressions, we need to truncate the summation to finite ranges of $\tilde{\Lambda} = \{\ell, m, n_r, n_\theta\}$ in Eqs. (26)-(28). In order to save the computation time in the numerical calculation, we sum ℓ up to 7. We can check that the error due to neglecting terms for $\ell \geq 8$ is smaller than the relative error in the 4PN $O(e^6)$ formulae from the corresponding numerical results up to $\ell = 7$. We also truncate n_r and n_θ to achieve the relative error of $\sim 10^{-7}$ in numerical results up to $\ell = 7$. For the parameters investigated in the comparison, the relative error of $\sim 10^{-7}$ achieved by truncating n_r and n_θ is again smaller than the relative error in the 4PN $O(e^6)$ formulae from the numerical results up to $\ell = 7$. Thus, we can regard numerical results as benchmarks to investigate the accuracy in our analytic formulae.

Here we define the relative error in the analytic formula of $\langle dE/dt \rangle_t$ by

$$\Delta_E \equiv \left| 1 - \left\langle \frac{dE}{dt} \right\rangle_t^{\text{Ana}} / \left\langle \frac{dE}{dt} \right\rangle_t^{\text{Num}} \right|, \quad (39)$$

where $\langle dE/dt \rangle_t^{\text{Ana}}$ denotes the analytic formula in order to distinguish it from the corresponding numerical result, $\langle dE/dt \rangle_t^{\text{Num}}$. We also define the relative errors in the analytic formulae of $\langle dL/dt \rangle_t$ and $\langle dC/dt \rangle_t$ in a similar manner and denote them as Δ_L and Δ_C respectively.

Fig. 1 shows several plots of Δ_E for the 4PN $O(e^6)$ formula as a function of p for several sets of (e, ι) with $q = 0.9$. In the plots, we also show the relative errors in the 2.5PN $O(e^2)$ and 3PN $O(e^4)$ formulae for reference. From the plots for $e = 0.1$ (three on the top), one can find that Δ_E falls off faster than p^{-4} for $p \gtrsim 10$ (Similarly, the relative errors in the 2.5PN $O(e^2)$ and 3PN $O(e^4)$ formulae fall off faster than $p^{-5/2}$ and p^{-3}). Noting $v = \sqrt{1/p}$, this would be a good indication that our PN formula has been derived correctly up to required order.

Δ_E is expected to contain not only higher order corrections than the 4PN order, but also the higher order corrections of eccentricity than $O(e^6)$ in the lower PN terms, which will become dominant when p and e get larger. In fact, seeing the plots for $e = 0.7$ in Fig. 1, one can find that the relative error strays out of the expected power law line for large p . This behavior is clearer in the plots of the relative error in the 2.5PN $O(e^2)$ formula. From Eqs. (32) and (36), we know that the relative error in the 2.5PN $O(e^2)$ formula contains the $O(e^4)$ correction in the $O(v^0)$ term. The effect of this correction appears as large- p plateaus in the plots (also see Fig. 6). This may motivate us to perform the higher order expansion with respect to the orbital eccentricity in the PN formulae or to derive the PN formulae without performing the expansion with respect to the orbital eccentricity [30–33]. In addition, it might be noted that the behavior of the relative error does not strongly depend on the inclination angle ι for fixed q and e .

In Fig. 2, we show the relative errors in the 4PN $O(e^6)$ formulae for the secular changes of the three orbital parameters, $\{E, L, C\}$, for several sets of (q, e) and $\iota = 50^\circ$. As in the case of Δ_E shown in Fig. 1, the relative errors, Δ_L and Δ_C , fall off faster than p^{-4} when $p \gtrsim 10$, except for the large p region ($p \gtrsim 100$) in the case of $e = 0.7$. Thus, the 4PN $O(e^6)$ formulae for the secular changes of the orbital parameters are expected to be valid up to $O(v^8)$. From Fig. 2, one might think that it is enough to investigate only Δ_E to discuss the accuracy of our formulae since there are not large differences in the relative errors, Δ_E , Δ_L and Δ_C .

Fig. 3 shows contour plots for Δ_E as a function of p and e for several sets of (ι, q) . From these plots, one may be able to comprehend the accuracy of our PN formulae more easily than using Figs. 1 and 2. One will find that the relative error becomes smaller (larger) for larger (smaller) p and smaller (larger) e . Moreover, it might be noticed that the relative error does not strongly depend on the inclination angle ι for fixed q as expected from Fig. 1. If one requires $\Delta_E < 10^{-5}$ as an error tolerance, one can use the contour line with the label 10^{-5} to find the region of validity in the figure. For example, one will find that $\Delta_E < 10^{-5}$ for $p \gtrsim 50$ and $e = 0.1$, $p \gtrsim 80$ and $e = 0.4$, and $p \gtrsim 120$ and $e = 0.7$ when $q = 0.9$.




Fig. 1 The relative errors in the analytic PN formulae for the secular change of the particle's energy as a function of the semi-latus rectum p for $q = 0.9$, $e = 0.1, 0.4$ and 0.7 (from top to bottom) and $\iota = 20^\circ, 50^\circ$ and 80° (from left to right). In addition to the error in the 4PN $O(e^6)$ formula, those in the 2.5PN $O(e^2)$ and the 3PN $O(e^4)$ formulae are shown in each plot for reference. We truncated the plots at $p = 6$ because the relative errors get too large (nearly or more than unity) in $p < 6$ to be meaningful. One finds that the relative error becomes smaller with increasing orders of the PN approximation and the expansion with respect to the eccentricity. The relative error in our 4PN $O(e^6)$ formula falls off faster than p^{-4} when the eccentricity is small, *e.g.* $e \lesssim 0.4$. Since $v = \sqrt{1/p}$, this would imply that our 4PN formula is correctly representing the secular change up to the 4PN order. Note, however, that the relative error in the 4PN $O(e^6)$ formula for $e = 0.7$ falls off slower than p^{-4} when the semi-latus rectum becomes larger, *e.g.* $p > 100$. This might be because of the higher order corrections of e than $O(e^6)$, which will contain the lower PN terms than the 4PN order. We also note that changing the inclination angle, ι , does not change the dependence on p of the relative error for fixed q and e so much. This might be checked more easily in contour plots in Fig. 3, which show the relative error as a function of p and e for fixed q and ι .

3.3. Implementation of an exponential resummation method

In order to improve the accuracy in the analytic PN formulae, one may apply some resummation methods such as Padé approximation [34], the factorized resummation [35–37] and the exponential resummation [38]. Since the exponential resummation may be the simplest




Fig. 2 The relative errors in the analytic PN formulae for the secular changes of the three orbital parameters, $\{E, L, C\}$, as functions of the semi-latus rectum p for $\iota = 50^\circ$, $q = 0.9, 0.5, 0.1$ and -0.9 (from top to bottom) and $e = 0.1, 0.4$ and 0.7 (from left to right). We truncated the plots at $p = \max\{6, p_s(e, \iota)\}$, where $p_s(e, \iota)$ is the value of p at the “separatrix” (the boundary between stable and unstable orbits), because the relative errors get too large in $p < 6$ to be meaningful and the orbit is not stable for $p < p_s(e, \iota)$. As pointed out in Fig. 1, the relative errors in our 4PN $O(e^6)$ formulae fall off faster than p^{-4} when the eccentricity is small, *e.g.* $e \lesssim 0.4$, while the fall-off gets slower when p is larger for $e = 0.7$. There are not large differences in the behaviors of Δ_E , Δ_L and Δ_C . This suggests that it might be enough to focus only on $\langle dE/dt \rangle_t$ to investigate the accuracy and convergence of our 4PN formulae.

one to implement among them, we here choose to implement the exponential resummation. We apply it to our 4PN formulae and check how the accuracy is improved.

To introduce the exponential resummation, we make use of the following identity

$$\left\langle \frac{dI}{dt} \right\rangle_t = \left(\frac{dI}{dt} \right)_N \exp \left\{ \ln \left[\left\langle \frac{dI}{dt} \right\rangle_t / \left(\frac{dI}{dt} \right)_N \right] \right\}, \quad (40)$$




Fig. 3 The relative error in the 4PN $O(e^6)$ formula for the secular change of the particle's energy, Δ_E , as a function of the semi-latus rectum p and the eccentricity e for $q = 0.9, 0.5, 0.1$ and -0.9 (from top to bottom) and $\iota = 20^\circ, 50^\circ$ and 80° (from left to right). We truncated the plots at $p = \max\{6, p_s(e, \iota)\}$ because the relative errors get too large in $p < 6$ to be meaningful and the orbit is not stable for $p < p_s(e, \iota)$. From the figures, it is easily found that the relative error becomes smaller (larger) for larger (smaller) p and smaller (larger) e with fixed q and ι . If one requires the relative error to be less than 10^{-5} , the region in the semi-latus rectum p and the eccentricity e will be $p \gtrsim 50$ and $e = 0.1$, $p \gtrsim 80$ and $e = 0.4$, and $p \gtrsim 120$ and $e = 0.7$ when $q = 0.9$. It might be noticed that the relative error does not strongly depend on the inclination angle ι for fixed q as pointed out in Fig. 1.

where $I = \{E, L, C\}$. The exponential resummation can be obtained by replacing the exponent in (40) to the expansion with respect to v ,

$$F_n^I := \ln \left[\left\langle \frac{dI}{dt} \right\rangle_t \middle/ \left(\frac{dI}{dt} \right)_N \right] \Big|_{\text{truncated after } n\text{th order of } v}, \quad (41)$$

where we do not perform the expansion with respect to e . Since our PN formulae for $\langle dI/dt \rangle_t$ are given at the 4PN order, we truncate F_n^I after $O(v^8)$. Finally, the exponential resummed form is expressed as

$$\left\langle \frac{dI}{dt} \right\rangle_t^{\text{exp}} = \left(\frac{dI}{dt} \right)_N \exp F_8^I. \quad (42)$$

Fig. 4 shows the relative errors in the exponential resummed forms of the secular changes of E , L and C , estimated by using Eq. (39). We also show the relative errors in the Taylor-type formulae in the same graphs for comparison. One will find that the relative errors in the exponential resummed forms are less than those in the Taylor-type formulae in most cases, except for $\langle dC/dt \rangle_t$ in the case of $q = 0.9$, $(e, \iota) = (0.1, 50^\circ)$. Using the exponential resummation when $q = 0.9$ and $\iota = 50^\circ$, the region to satisfy $\Delta_E < 10^{-5}$ is extended to $p \gtrsim 40$ from $p \gtrsim 50$ for $e = 0.1$, $p \gtrsim 60$ from $p \gtrsim 80$ for $e = 0.4$, and $p \gtrsim 100$ from $p \gtrsim 120$ for $e = 0.7$. This might motivate us to use the resummation method to improve the accuracy of Taylor-type formulae even in the case of general orbits.

3.4. Convergence with respect to v and e of the analytic formulae

Apart from comparisons to the numerical results, we may also discuss the convergence property in our PN formulae with respect to v and e by investigating the contribution of each order of v and e in the formulae although this is a rough estimation.

First we assess the PN convergence of our formulae. For this purpose, we introduce Δ_n as

$$\left\langle \frac{dE}{dt} \right\rangle_t^{\text{PN}} = \left(\frac{dE}{dt} \right)_N \sum_{n=0}^8 \Delta_n, \quad (43)$$

where $p = 1/v^2$ and Δ_n is the $O(v^n)$ term in the PN formula of $\langle dE/dt \rangle_t$, *e.g.* $\Delta_0 = 1 + \frac{73}{24}e^2 + \frac{37}{96}e^4$, $\Delta_1 = 0$ and $\Delta_2 = \left(-\frac{1247}{336} - \frac{9181}{672}e^2 + \frac{809}{128}e^4 + \frac{8609}{5376}e^6\right)v^2$. Δ_n depends on (q, p, e, Y) in general although we omit the argument for simplicity.

Since Δ_n shows the relative importance of the $O(v^n)$ term in the PN formulae, it can be used to investigate the convergence property with respect to v : it is expected that $|\Delta_{n+1}| < |\Delta_n|$ for moderately large n if the PN formula converges. In Fig. 5, we plot the relative contribution of each order, Δ_n , as a function of p for several sets of (e, ι) and $q = 0.9$. From this figure, one may find that Δ_n does not strongly depend on the inclination angle, ι , as shown in Sec. 3.2, while it strongly depends on e . The convergence gets worse when the orbital eccentricity becomes larger. This tendency is particularly evident in the small- p region. Fixing the value of p , the orbit with larger e passes by closer to the central black hole and will be affected by the stronger gravitational field. Hence the PN convergence is expected to be worse when the eccentricity becomes larger.

Next, in order to investigate the convergence of the expansion with respect to the orbital eccentricity in the PN formula, we introduce A_n as

$$\left\langle \frac{dE}{dt} \right\rangle_t^{\text{PN}} = \left(\frac{dE}{dt} \right)_N [A_0 e^0 + A_2 e^2 + A_4 e^4 + A_6 e^6], \quad (44)$$

where the term A_0 coincides with the energy flux for circular orbits and $A_n = 0$ when n is odd.

One may ask whether the condition, $|A_{2n+2}e^{2n+2}| < |A_{2n}e^{2n}|$, is satisfied for moderately large integer n if the series converges. From Fig. 6, it is found that the condition is satisfied




Fig. 4 The relative errors in the PN formulae and the exponential resummation formulae for the secular changes of the orbital parameters, $\{E, L, C\}$ as functions of the semi-latus rectum p for $q = 0.9$, $\iota = 50^\circ$ and $e = 0.1, 0.4$ and 0.7 (from top to bottom). We truncated the plots at $p = 6$ because the relative errors in the PN formulae get too large in $p < 6$ to be meaningful. Using the exponential resummation, the accuracy is improved in most cases. For example, the region to satisfy $\Delta_E < 10^{-5}$ is improved from $p \gtrsim 50$ to $p \gtrsim 40$ for $e = 0.1$, $p \gtrsim 80$ to $p \gtrsim 60$ for $e = 0.4$, and $p \gtrsim 120$ to $p \gtrsim 100$ for $e = 0.7$. This would suggest us to try to apply resummation methods to the PN formulae even in the case of general orbits.

in most cases. As expected, the convergence becomes slower when the eccentricity is larger. Especially, the convergence gets worse when $p \lesssim 10$ in $e = 0.7$ case. The calculation of the higher PN corrections will be necessary to improve the bad convergence for small p . We also note that A_n does not strongly depend on ι for a fixed q as in Sec. 3.2.

4. Summary

We have derived the secular changes of the orbital parameters, the energy, azimuthal angular momentum, and Carter parameter of a point particle orbiting a Kerr black hole, by using the post-Newtonian approximation in the first order black hole perturbation theory. We have extended the previous work [15], which derived formulae up to the 2.5PN order with the second order correction with respect to the eccentricity, to the 4PN order with the sixth order correction with respect to the eccentricity. We have also included the contribution due to the black hole absorption, which has not been included in [15]. As shown in the case of equatorial, circular orbits [27, 28], we have found that the secular changes of the




Fig. 5 The relative contribution of the $O(v^n)$ term in the PN formula for $\langle dE/dt \rangle_t$, defined in Eq. (43). We plot the absolute value of Δ_n as a function of the semi-latus rectum p for $e = 0.1, 0.4$ and 0.7 (from left to right), and $\iota = 20^\circ, 50^\circ$ and 80° (from top to bottom) when $q = 0.9$. We truncated the plots at $p = 6$ because the relative contributions get too large in $p < 6$ to be meaningful. It is expected that $|\Delta_{n+1}| < |\Delta_n|$ for moderately large n if the PN formula converges. As shown in Sec. 3.2, Δ_n does not strongly depend on ι for a fixed e although Δ_n strongly depends on e . In fact, the convergence seems worse the orbital eccentricity becomes larger. This tendency is clear for small p , *e.g.* $p \lesssim 10$.

three orbital parameters due to the absorption, appear at the 2.5PN (4PN) from the leading order in the Kerr (Schwarzschild) case, and that the 2.5PN and 3.5PN contributions of the absorption to the secular change of the particle's energy can be positive for $q > 0$, which implies that a superradiance can be realized in the Kerr case. We have also found that the superradiant contributions in the secular change of the energy get smaller when the inclination angle becomes larger and they vanishes for polar ($Y = 0$) orbits. This means that the superradiant scattering may be suppressed for inclined orbits [29].

To investigate the accuracy in our 4PN formulae, we have compared the formulae to high-precision numerical results [18] in Sec. 3.2. We have found that the accuracy gets worse when the orbital velocity and the orbital eccentricity become larger, as expected. If the relative error in the 4PN $O(e^6)$ formula for the secular change of the energy is required to be less than 10^{-5} , the parameter region to satisfy it might be $p \gtrsim 50$ for $e = 0.1$, $p \gtrsim 80$ for $e = 0.4$, and $p \gtrsim 120$ for $e = 0.7$ when $q = 0.9$. The region does not strongly depend on the orbital inclination angle. From Fig. 1, one can clearly find the improvement of the accuracy in our




Fig. 6 The relative contribution of the $O(e^n)$ term in the PN formula for $\langle dE/dt \rangle_t$, defined in Eq. (44). We plot the absolute value of $A_n e^n$ as a function of the semi-latus rectum p for $\iota = 20^\circ, 50^\circ$ and 80° (from left to right) when $q = 0.9$. We truncated the plots at $p = 6$ because the relative contributions get too large in $p < 6$ to be meaningful. It is expected that $|A_{2n+2}e^{2n+2}| < |A_{2n}e^{2n}|$ for moderately large n if the series with respect to e converges. This condition is satisfied in most cases shown in this figure. The convergence becomes slower when the eccentricity is larger. Especially, the convergence for $p \lesssim 10$ is quite bad in $e = 0.7$ case. We also note that A_n does not strongly depend on ι for a fixed q as mentioned in Sec. 3.2.

PN formulae from the previous work at the 2.5PN order and the second order correction in the orbital eccentricity [15] whose relative error is larger than 10^{-2} for $p \gtrsim 100$ when $e \gtrsim 0.4$ since, in this region, the error due to the truncation of the expansion with respect to the orbital eccentricity is larger than the one of the PN expansion.

One may improve the accuracy of our PN formulae by using resummation methods. In this paper, we have applied the exponential resummation [38] to our 4PN formulae and confirmed that the resummation method improves the accuracy in most cases investigated here. For example, we found that the region in which the relative errors are less than 10^{-5} can be extended from $p \gtrsim 50$ to $p \gtrsim 40$ for $e = 0.1$, $p \gtrsim 80$ to $p \gtrsim 60$ for $e = 0.4$, and $p \gtrsim 120$ to $p \gtrsim 100$ for $e = 0.7$.

We also investigate the convergence properties of the PN expansion and the expansion with respect to the orbital eccentricity, respectively. Both convergences get worse when the semi-latus rectum is smaller; in other words, the gravitational field becomes stronger. This

tendency gets clearer in the case of large eccentricity, in which the particle passes by closer to the central black hole.

In order to improve the accuracy and convergence of the 4PN $O(e^6)$ formulae near the central black hole and to obtain the physical information of the source in the strong-field region, it is necessary to derive the higher order corrections of the PN expansion and the expansion with respect to the eccentricity. It may be possible to avoid the expansion with respect to the eccentricity and to derive the PN formulae applicable to arbitrary eccentricity. So far the PN formulae of the rate of the energy loss without performing the expansion with respect to the eccentricity had been derived for equatorial orbits in [30–33]. The extension of these results to the case of inclined orbits is challenging: we can obtain analytic expressions for general bound geodesic orbits in Kerr spacetime without performing the expansion with respect to the eccentricity nor the inclination by using results in Ref. [39], while we need to reformulate the source term of the Teukolsky equation and the derivation of the partial waves constructed from the source term. We would like to leave it to the future work.

Acknowledgment

We would like to thank Takahiro Tanaka and Hiroyuki Nakano for useful discussions and comments. We are also grateful to Theoretical Astrophysics Group in Kyoto University for hospitality during the intermediate stage of completing this paper. NS acknowledges the support of the Grand-in-Aid for Scientific Research (No. 25800154). RF’s work was supported by the European Union’s FP7 ERC Starting Grant “The dynamics of black holes: testing the limits of Einstein’s theory” grant agreement no. DyBHo-256667. Some numerical computations were performed at the cluster “Baltasar-Sete-Sóis” in CENTRA/IST. Some analytic calculations were carried out on HA8000/RS440 at Yukawa Institute for Theoretical Physics in Kyoto University.

A. PN formulae for the orbital parameters and fundamental frequencies

In this section, we present the PN formulae of the orbital parameters, $\{\hat{E}, \hat{L}, \hat{C}\}$, and the fundamental frequencies, $\{\Omega_t, \Omega_r, \Omega_\theta, \Omega_\phi\}$. Here we show the formulae up to the 3PN $O(e^6)$ order to save space although it is possible to calculate them to the higher order. The higher order results will be publicly available online [26].

$$\begin{aligned}
\hat{E} &= 1 + \left\{ -\frac{1}{2} + \frac{1}{2} e^2 \right\} v^2 + \left\{ \frac{3}{8} - \frac{3}{4} e^2 + \frac{3}{8} e^4 \right\} v^4 + \{ -Yq + 2Yqe^2 - Yqe^4 \} v^5 \\
&\quad + \left\{ \frac{27}{16} + \frac{1}{2} Y^2 q^2 + \left(-\frac{49}{16} - Y^2 q^2 \right) e^2 + \left(\frac{17}{16} + \frac{1}{2} Y^2 q^2 \right) e^4 + \frac{5}{16} e^6 \right\} v^6, \quad (\text{A1}) \\
\hat{L} &= pv \left[Y + \left\{ \frac{3}{2} Y + \frac{1}{2} Y e^2 \right\} v^2 + \{ -3Y^2 q - qY^2 e^2 \} v^3 \right. \\
&\quad + \left\{ \frac{27}{8} Y + Y^3 q^2 + \left(\frac{9}{4} Y + Y^3 q^2 \right) e^2 + \frac{3}{8} Y e^4 \right\} v^4 \\
&\quad + \left\{ -\frac{15}{2} Y^2 q - 7qY^2 e^2 - \frac{3}{2} Y^2 q e^4 \right\} v^5 \\
&\quad \left. + \left\{ \frac{135}{16} Y - 4Yq^2 + 9Y^3 q^2 + \left(\frac{135}{16} Y - 4Yq^2 + 9Y^3 q^2 \right) e^2 \right\} v^6 \right]
\end{aligned}$$

$$+ \left(\frac{45}{16} Y + 2 Y^3 q^2 \right) e^4 + \frac{5}{16} Y e^6 \Big\} v^6 \Big], \quad (\text{A2})$$

$$\begin{aligned} \hat{C} &= \left\{ \frac{1}{Y^2} - 1 \right\} L^2, \\ &= p^2 v^2 (1 - Y^2) \Big[1 + (3 + e^2) v^2 - 2 q Y (3 + e^2) v^3 \\ &\quad + \{ 9 + 2 Y^2 q^2 + (6 + 2 Y^2 q^2) e^2 + e^4 \} v^4 - 4 q Y (2 + e^2) (3 + e^2) v^5 \\ &\quad + \{ 27 - 8 q^2 + 30 Y^2 q^2 + (27 + 28 Y^2 q^2 - 8 q^2) e^2 + (9 + 6 Y^2 q^2) e^4 + e^6 \} v^6 \Big], \quad (\text{A3}) \end{aligned}$$

$$\begin{aligned} \Omega_t &= p^2 \Big[1 + \frac{3}{2} e^2 + \frac{15}{8} e^4 + \frac{35}{16} e^6 + \left\{ \frac{3}{2} - \frac{1}{4} e^2 - \frac{15}{16} e^4 - \frac{45}{32} e^6 \right\} v^2 \\ &\quad + \left\{ 2 Y q e^2 + 3 Y q e^4 + \frac{15}{4} Y q e^6 \right\} v^3 \\ &\quad + \left\{ \frac{27}{8} - \frac{1}{2} Y^2 q^2 + \frac{1}{2} q^2 + \left(-\frac{99}{16} + q^2 - 2 Y^2 q^2 \right) e^2 \right. \\ &\quad \left. + \left(-\frac{567}{64} + \frac{21}{16} q^2 - \frac{45}{16} Y^2 q^2 \right) e^4 + \left(-\frac{1371}{128} + \frac{25}{16} q^2 - \frac{55}{16} Y^2 q^2 \right) e^6 \right\} v^4 \\ &\quad + \left\{ -3 Y q + \frac{43}{2} Y q e^2 + \frac{231}{8} Y q e^4 + \frac{555}{16} Y q e^6 \right\} v^5 \\ &\quad + \left\{ \frac{135}{16} - \frac{1}{4} q^2 + \frac{3}{4} Y^2 q^2 + \left(-\frac{1233}{32} + \frac{47}{4} q^2 - \frac{75}{2} Y^2 q^2 \right) e^2 \right. \\ &\quad \left. + \left(-\frac{6567}{128} + \frac{499}{32} q^2 - \frac{1577}{32} Y^2 q^2 \right) e^4 \right. \\ &\quad \left. + \left(-\frac{15565}{256} + \frac{75}{4} q^2 - \frac{1887}{32} Y^2 q^2 \right) e^6 \right\} v^6 \Big], \quad (\text{A4}) \end{aligned}$$

$$\begin{aligned} \Omega_r &= p v \Big[1 + \left\{ -\frac{3}{2} + \frac{1}{2} e^2 \right\} v^2 + \{ 3 Y q - Y q e^2 \} v^3 \\ &\quad + \left\{ -\frac{45}{8} + \frac{1}{2} q^2 - 2 Y^2 q^2 + \left(\frac{1}{4} q^2 + \frac{1}{4} Y^2 q^2 \right) e^2 + \frac{3}{8} e^4 \right\} v^4 \\ &\quad + \left\{ \frac{33}{2} Y q + 2 Y q e^2 - \frac{3}{2} Y q e^4 \right\} v^5 \\ &\quad + \left\{ -\frac{351}{16} - \frac{51}{2} Y^2 q^2 + \frac{33}{4} q^2 + \left(-\frac{135}{16} + \frac{7}{8} q^2 - \frac{39}{8} Y^2 q^2 \right) e^2 \right. \\ &\quad \left. + \left(\frac{21}{16} + \frac{1}{8} q^2 + \frac{13}{8} Y^2 q^2 \right) e^4 + \frac{5}{16} e^6 \right\} v^6 \Big], \quad (\text{A5}) \end{aligned}$$

$$\begin{aligned} \Omega_\theta &= p v \Big[1 + \left\{ \frac{3}{2} + \frac{1}{2} e^2 \right\} v^2 - \{ 3 Y q + Y q e^2 \} v^3 \\ &\quad + \left\{ \frac{27}{8} + \frac{7}{4} Y^2 q^2 - \frac{1}{4} q^2 + \left(\frac{9}{4} + \frac{1}{4} q^2 + \frac{1}{4} Y^2 q^2 \right) e^2 + \frac{3}{8} e^4 \right\} v^4 \\ &\quad - \left\{ \frac{15}{2} Y q + 7 Y q e^2 + \frac{3}{2} Y q e^4 \right\} v^5 \end{aligned}$$

$$\begin{aligned}
& + \left\{ \frac{135}{16} + \frac{57}{8} Y^2 q^2 - \frac{27}{8} q^2 + \left(\frac{135}{16} - \frac{19}{4} q^2 + \frac{45}{4} Y^2 q^2 \right) e^2 \right. \\
& \quad \left. + \left(\frac{45}{16} + \frac{1}{8} q^2 + \frac{13}{8} Y^2 q^2 \right) e^4 + \frac{5}{16} e^6 \right\} v^6 \Big], \tag{A6}
\end{aligned}$$

$$\begin{aligned}
\Omega_\varphi &= p v \left[1 + \left\{ \frac{3}{2} + \frac{1}{2} e^2 \right\} v^2 + \{ 2q - 3Yq - Yq e^2 \} v^3 \right. \\
& + \left\{ -\frac{3}{2} Yq^2 + \frac{7}{4} Y^2 q^2 - \frac{1}{4} q^2 + \frac{27}{8} + \left(\frac{9}{4} + \frac{1}{4} q^2 + \frac{1}{4} Y^2 q^2 \right) e^2 + \frac{3}{8} e^4 \right\} v^4 \\
& + \left\{ 3q - \frac{15}{2} Yq + (4q - 7Yq) e^2 - \frac{3}{2} Yq e^4 \right\} v^5 \\
& + \left\{ -\frac{9}{4} Yq^2 + \frac{57}{8} Y^2 q^2 + \frac{135}{16} - \frac{27}{8} q^2 + \left(\frac{135}{16} - \frac{19}{4} q^2 - \frac{35}{4} Yq^2 + \frac{45}{4} Y^2 q^2 \right) e^2 \right. \\
& \quad \left. + \left(\frac{45}{16} + \frac{1}{8} q^2 + \frac{13}{8} Y^2 q^2 \right) e^4 + \frac{5}{16} e^6 \right\} v^6 \Big]. \tag{A7}
\end{aligned}$$

B. Fourier coefficients of bound orbits

Here we show the PN formulae of the Fourier coefficients in Eqs. (15), (16), (18) and (19) up to the 3PN $O(e^6)$ order. The 4PN $O(e^6)$ results obtained in this work will be available online [26].

In this work, we follow the same procedure as in [15] to derive the amplitudes of the partial waves, $Z_\Lambda^{\text{H},\infty}$ in (23). In the formal expression, the dependence of $\varphi^{(\theta)}$ appears in the form of the combination as $X \equiv \sin \theta e^{i\varphi^{(\theta)}}$, which can be expressed in the Fourier series as

$$X = p \sum_{n_\theta=0}^{\infty} \left[X_{n_\theta}^{\Re} \cos n_\theta \Omega_\theta \lambda + i X_{n_\theta}^{\Im} \sin n_\theta \Omega_\theta \lambda \right], \tag{B1}$$

Therefore we show the Fourier coefficients of X instead of $\varphi^{(\theta)}$.

B.1. Radial component

$$\begin{aligned}
\alpha_0 &= 1 + e^2 \left\{ \frac{1}{2} - \frac{1}{2} v^2 + qYv^3 + \left(-3 + \left(\frac{1}{2} - Y^2 \right) q^2 \right) v^4 \right. \\
& \quad \left. + 10qYv^5 + \left(-18 + \left(\frac{11}{2} - 18Y^2 \right) q^2 \right) v^6 \right\} \\
& + e^4 \left\{ \frac{3}{8} - \frac{3}{8} v^2 + \frac{3}{4} qYv^3 + \left(-\frac{33}{16} + \left(\frac{5}{16} - \frac{11}{16} Y^2 \right) q^2 \right) v^4 \right. \\
& \quad \left. + \frac{27}{4} qYv^5 + \left(-\frac{189}{16} + \left(-\frac{185}{16} Y^2 + \frac{61}{16} \right) q^2 \right) v^6 \right\} \\
& + e^6 \left\{ \frac{5}{16} - \frac{5}{16} v^2 + \frac{5}{8} qYv^3 + \left(-\frac{27}{16} + \left(\frac{1}{4} - \frac{9}{16} Y^2 \right) q^2 \right) v^4 \right. \\
& \quad \left. + \frac{11}{2} qYv^5 + \left(-\frac{19}{2} + \left(-\frac{75}{8} Y^2 + \frac{49}{16} \right) q^2 \right) v^6 \right\}, \tag{B2} \\
\alpha_1 &= e + e^3 \left\{ \frac{3}{4} - \frac{1}{2} v^2 + Yqv^3 + \left(-\frac{51}{16} + \left(\frac{7}{16} - \frac{15}{16} Y^2 \right) q^2 \right) v^4 \right.
\end{aligned}$$

$$\begin{aligned}
& +\frac{43}{4}Yqv^5 + \left(-\frac{81}{4} + \left(-\frac{153}{8}Y^2 + \frac{11}{2}\right)q^2\right)v^6\Big\} \\
& +e^5\left\{\frac{5}{8} - \frac{1}{2}v^2 + Yqv^3 + \left(-\frac{93}{32} + \left(\frac{13}{32} - \frac{29}{32}Y^2\right)q^2\right)v^4\right. \\
& \left. +\frac{77}{8}Yqv^5 + \left(-\frac{277}{16} + \left(-\frac{133}{8}Y^2 + \frac{83}{16}\right)q^2\right)v^6\right\}, \tag{B3}
\end{aligned}$$

$$\begin{aligned}
\alpha_2 = & e^2\left\{\frac{1}{2} + \frac{1}{2}v^2 - Yqv^3 + \left(3 + \left(Y^2 - \frac{1}{2}\right)q^2\right)v^4\right. \\
& \left.-10Yqv^5 + \left(18 + \left(18Y^2 - \frac{11}{2}\right)q^2\right)v^6\right\} \\
& +e^4\left\{\frac{1}{2} - \frac{1}{2}v^4 + 2Yqv^5 + \left(-\frac{11}{2} + \left(-4Y^2 + \frac{1}{2}\right)q^2\right)v^6\right\} \\
& +e^6\left\{\frac{15}{32} - \frac{5}{32}v^2 + \frac{5}{16}Yqv^3 + \left(-\frac{39}{32} + \left(\frac{1}{8} - \frac{9}{32}Y^2\right)q^2\right)v^4\right. \\
& \left. +\frac{17}{4}Yqv^5 + \left(-\frac{279}{32} + \left(-\frac{243}{32}Y^2 + 2\right)q^2\right)v^6\right\}, \tag{B4}
\end{aligned}$$

$$\begin{aligned}
\alpha_3 = & e^3\left\{\frac{1}{4} + \frac{v^2}{2} - Yqv^3 + \left(\frac{51}{16} + \left(-\frac{7}{16} + \frac{15}{16}Y^2\right)q^2\right)v^4\right. \\
& \left.-\frac{43}{4}Yqv^5 + \left(\frac{81}{4} + \left(\frac{153}{8}Y^2 - \frac{11}{2}\right)q^2\right)v^6\right\} \\
& +e^5\left\{\frac{5}{16} + \frac{v^2}{4} - \frac{Yqv^3}{2} + \left(\frac{69}{64} + \left(-\frac{13}{64} + \frac{29}{64}Y^2\right)q^2\right)v^4\right. \\
& \left.-\frac{53}{16}Yqv^5 + \left(\frac{135}{32} + \left(\frac{43}{8}Y^2 - \frac{69}{32}\right)q^2\right)v^6\right\}, \tag{B5}
\end{aligned}$$

$$\begin{aligned}
\alpha_4 = & e^4\left\{\frac{1}{8} + \frac{3}{8}v^2 - \frac{3}{4}Yqv^3 + \left(\frac{41}{16} + \left(-\frac{5}{16} + \frac{11}{16}Y^2\right)q^2\right)v^4\right. \\
& \left.-\frac{35}{4}Yqv^5 + \left(\frac{277}{16} + \left(\frac{249}{16}Y^2 - \frac{69}{16}\right)q^2\right)v^6\right\} \\
& +e^6\left\{\frac{3}{16} + \frac{5}{16}v^2 - \frac{5}{8}Yqv^3 + \left(\frac{27}{16} + \left(-\frac{1}{4} + \frac{9}{16}Y^2\right)q^2\right)v^4\right. \\
& \left.-\frac{11}{2}Yqv^5 + \left(9 + \left(\frac{75}{8}Y^2 - \frac{49}{16}\right)q^2\right)v^6\right\}, \tag{B6}
\end{aligned}$$

$$\begin{aligned}
\alpha_5 = & e^5\left\{\frac{1}{16} + \frac{v^2}{4} - \frac{Yq}{2}v^3 + \left(\frac{117}{64} + \left(-\frac{13}{64} + \frac{29}{64}Y^2\right)q^2\right)v^4\right. \\
& \left.-\frac{101}{16}Yqv^5 + \left(\frac{419}{32} + \left(\frac{45}{4}Y^2 - \frac{97}{32}\right)q^2\right)v^6\right\}, \tag{B7}
\end{aligned}$$

$$\begin{aligned}
\alpha_6 = & e^6\left\{\frac{1}{32} + \frac{5}{32}v^2 - \frac{5}{16}Yqv^3 + \left(\frac{39}{32} + \left(-\frac{1}{8} + \frac{9}{32}Y^2\right)q^2\right)v^4\right. \\
& \left.-\frac{17}{4}Yqv^5 + \left(\frac{295}{32} + \left(\frac{243}{32}Y^2 - 2\right)q^2\right)v^6\right\}, \tag{B8}
\end{aligned}$$

$$\alpha_n = O(e^n) \quad \text{for } n \geq 7. \tag{B9}$$

B.2. Longitudinal component

$$\beta_0 = 0, \quad (B10)$$

$$\begin{aligned} \beta_1 = & 1 + \left(\frac{1}{16} - \frac{9}{16} Y^2 \right) q^2 v^4 + \left(-\frac{1}{4} + \frac{9}{4} Y^2 \right) q^2 v^6 \\ & + e^2 \left\{ \left(-\frac{1}{16} + \frac{9}{16} Y^2 \right) q^2 v^4 + \left(-\frac{9}{4} Y^2 + \frac{1}{4} \right) q^2 v^6 \right\}, \end{aligned} \quad (B11)$$

$$\beta_2 = 0, \quad (B12)$$

$$\beta_3 = \frac{1-Y^2}{16} q^2 v^4 - \frac{1-Y^2}{4} q^2 v^6 + e^2 \left\{ -\frac{1-Y^2}{16} q^2 v^4 + \frac{1-Y^2}{4} q^2 v^6 \right\}, \quad (B13)$$

$$\beta_n = \begin{cases} 0 & (n: \text{even}) \\ O(v^{2n-2}) & (n: \text{odd}) \end{cases} \quad (B14)$$

B.3. r-part of the temporal component

$$\begin{aligned} \frac{v}{p} \tilde{t}_1^{(r)} = & e \left\{ 2 + 4v^2 - 6Yqv^3 + (17 + (4Y^2 - 1)q^2)v^4 - 54Yqv^5 \right. \\ & + (88 + (84Y^2 - 20)q^2)v^6 \Big\} \\ & + e^3 \left\{ 3 + 3v^2 - 4Yqv^3 + \left(\frac{77}{8} + \left(\frac{21}{8}Y^2 - \frac{5}{8} \right) q^2 \right) v^4 - \frac{57}{2}Yqv^5 \right. \\ & + \left(\frac{173}{4} + \left(42Y^2 - \frac{51}{4} \right) q^2 \right) v^6 \Big\} \\ & + e^5 \left\{ \frac{15}{4} + \frac{5}{2}v^2 - \frac{13}{4}Yqv^3 + \left(\frac{15}{2} + \left(\frac{17}{8}Y^2 - \frac{1}{2} \right) q^2 \right) v^4 - \frac{45}{2}Yqv^5 \right. \\ & + \left(\frac{67}{2} + \left(\frac{133}{4}Y^2 - 10 \right) q^2 \right) v^6 \Big\}, \end{aligned} \quad (B15)$$

$$\begin{aligned} \frac{v}{p} \tilde{t}_2^{(r)} = & e^2 \left\{ \frac{3}{4} + \frac{7}{4}v^2 - \frac{13}{4}Yqv^3 + \left(\frac{81}{8} + \left(5/2Y^2 - \frac{7}{8} \right) q^2 \right) v^4 - \frac{135}{4}Yqv^5 \right. \\ & + \left(\frac{499}{8} + \left(55Y^2 - \frac{113}{8} \right) q^2 \right) v^6 \Big\} \\ & + e^4 \left\{ \frac{5}{4} + \frac{7}{4}v^2 - 3Yqv^3 + \left(\frac{131}{16} + \left(\frac{37}{16}Y^2 - \frac{13}{16} \right) q^2 \right) v^4 - \frac{103}{4}Yqv^5 \right. \\ & + \left(\frac{691}{16} + \left(\frac{655}{16}Y^2 - \frac{197}{16} \right) q^2 \right) v^6 \Big\} \\ & + e^6 \left\{ \frac{105}{64} + \frac{105}{64}v^2 - \frac{175}{64}Yqv^3 + \left(\frac{905}{128} + \left(\frac{135}{64}Y^2 - \frac{95}{128} \right) q^2 \right) v^4 - \frac{1413}{64}Yqv^5 \right. \\ & + \left(\frac{4591}{128} + \left(\frac{2241}{64}Y^2 - \frac{1389}{128} \right) q^2 \right) v^6 \Big\}, \end{aligned} \quad (B16)$$

$$\begin{aligned} \frac{v}{p} \tilde{t}_3^{(r)} = & e^3 \left\{ \frac{1}{3} + v^2 - 2Yqv^3 + \left(\frac{53}{8} + \left(\frac{13}{8}Y^2 - \frac{5}{8} \right) q^2 \right) v^4 - \frac{45}{2}Yqv^5 \right. \\ & + \left(\frac{523}{12} + \left(38Y^2 - \frac{39}{4} \right) q^2 \right) v^6 \Big\} \end{aligned}$$

$$\begin{aligned}
& +e^5 \left\{ \frac{5}{8} + \frac{5}{4}v^2 - \frac{19}{8}Yqv^3 + \left(7 + \left(\frac{31}{16}Y^2 - \frac{3}{4} \right) q^2 \right) v^4 - \frac{91}{4}Yqv^5 \right. \\
& \left. + \left(\frac{647}{16} + \left(\frac{601}{16}Y^2 - \frac{175}{16} \right) q^2 \right) v^6 \right\}, \tag{B17}
\end{aligned}$$

$$\begin{aligned}
\frac{v}{p} \tilde{t}_4^{(r)} &= e^4 \left\{ \frac{5}{32} + \frac{19}{32}v^2 - \frac{39}{32}Yqv^3 + \left(\frac{137}{32} + \left(\frac{65}{64}Y^2 - \frac{13}{32} \right) q^2 \right) v^4 - \frac{473}{32}Yqv^5 \right. \\
& \left. + \left(\frac{957}{32} + \left(\frac{1631}{64}Y^2 - \frac{207}{32} \right) q^2 \right) v^6 \right\} \\
& + e^6 \left\{ \frac{21}{64} + \frac{57}{64}v^2 - \frac{113}{64}Yqv^3 + \left(\frac{89}{16} + \left(\frac{189}{128}Y^2 - \frac{19}{32} \right) q^2 \right) v^4 - \frac{1185}{64}Yqv^5 \right. \\
& \left. + \left(\frac{553}{16} + \left(\frac{4005}{128}Y^2 - \frac{141}{16} \right) q^2 \right) v^6 \right\}, \tag{B18}
\end{aligned}$$

$$\begin{aligned}
\frac{v}{p} \tilde{t}_5^{(r)} &= e^5 \left\{ \frac{3}{40} + \frac{7}{20}v^2 - \frac{29}{40}Yqv^3 + \left(\frac{27}{10} + \left(\frac{49}{80}Y^2 - 1/4 \right) q^2 \right) v^4 - \frac{189}{20}Yqv^5 \right. \\
& \left. + \left(\frac{319}{16} + \left(\frac{1321}{80}Y^2 - \frac{331}{80} \right) q^2 \right) v^6 \right\}, \tag{B19}
\end{aligned}$$

$$\begin{aligned}
\frac{v}{p} \tilde{t}_6^{(r)} &= e^6 \left\{ \frac{7}{192} + \frac{13}{64}v^2 - \frac{27}{64}Yqv^3 + \left(\frac{213}{128} + \left(\frac{23}{64}Y^2 - \frac{19}{128} \right) q^2 \right) v^4 - \frac{377}{64}Yqv^5 \right. \\
& \left. + \left(\frac{4969}{384} + \left(\frac{665}{64}Y^2 - \frac{329}{128} \right) q^2 \right) v^6 \right\}, \tag{B20}
\end{aligned}$$

$$\frac{v}{p} \tilde{t}_n^{(r)} = O(e^n) \quad (\text{for } n \geq 7). \tag{B21}$$

B.4. θ -part of the temporal component

$$\frac{1}{p} \tilde{t}_1^{(\theta)} = 0, \tag{B22}$$

$$\frac{1}{p} \tilde{t}_2^{(\theta)} = \frac{(Y^2 - 1)}{4} q^2 v^3 - \frac{(Y^2 - 1)}{2} q^2 v^5 + \frac{Y(Y^2 - 1)}{4} (3 + e^2) q^3 v^6, \tag{B23}$$

$$\frac{1}{p} \tilde{t}_n^{(\theta)} = \begin{cases} 0 & (n: \text{odd}) \\ O(v^{2n-1}) & (n: \text{even}) \end{cases} \tag{B24}$$

B.5. r -part of the azimuthal component

$$\tilde{\varphi}_1^{(r)} = e \left\{ -2qv^3 + 2Yq^2v^4 - 10qv^5 + 18Yq^2v^6 \right\}, \tag{B25}$$

$$\tilde{\varphi}_2^{(r)} = e^2 \left\{ -\frac{1}{4}Yq^2v^4 + \frac{1}{2}qv^5 - \frac{3}{4}Yq^2v^6 \right\}, \tag{B26}$$

$$\tilde{\varphi}_n^{(r)} = \begin{cases} O(v^{2n+1}) & (n: \text{odd}) \\ O(v^{2n}) & (n: \text{even}) \end{cases} \tag{B27}$$

B.6. θ -part of the azimuthal component

$$X_0^{\Re} = \left\{ \frac{1+Y}{2} - \frac{(9Y-1)(Y^2-1)}{32} q^2 v^4 + \frac{(9Y-1)(Y^2-1)}{8} q^2 v^6 \right\}$$

$$+e^2 \left\{ \frac{(9Y-1)(Y^2-1)}{32} q^2 v^4 - \frac{(9Y-1)(Y^2-1)}{8} q^2 v^6 \right\}, \quad (\text{B28})$$

$$X_1^{\mathfrak{R}} = 0, \quad (\text{B29})$$

$$X_2^{\mathfrak{R}} = \left\{ \frac{1-Y}{2} + \frac{Y(Y^2-1)}{4} q^2 v^4 - Y(Y^2-1) q^2 v^6 \right\} \\ +e^2 \left\{ -\frac{Y(Y^2-1)}{4} q^2 v^4 + Y(Y^2-1) q^2 v^6 \right\}, \quad (\text{B30})$$

$$X_3^{\mathfrak{R}} = 0, \quad (\text{B31})$$

$$X_4^{\mathfrak{R}} = \left\{ \frac{(Y+1)(Y-1)^2}{32} q^2 v^4 - \frac{(Y+1)(Y-1)^2}{8} q^2 v^6 \right\} \\ +e^2 \left\{ -\frac{(Y+1)(Y-1)^2}{32} q^2 v^4 + \frac{(Y+1)(Y-1)^2}{8} q^2 v^6 \right\}, \quad (\text{B32})$$

$$X_n^{\mathfrak{R}} = \begin{cases} 0 & (n: \text{odd}) \\ O(v^{2n-4}) & (n \geq 6: \text{even}) \end{cases}, \quad (\text{B33})$$

$$X_0^{\mathfrak{S}} = 0, \quad (\text{B34})$$

$$X_1^{\mathfrak{S}} = 0, \quad (\text{B35})$$

$$X_2^{\mathfrak{S}} = \left\{ \frac{Y-1}{2} - \frac{(5Y+1)(Y^2-1)}{16} q^2 v^4 + \frac{(5Y+1)(Y^2-1)}{4} q^2 v^6 \right\} \\ +e^2 \left\{ \frac{(5Y+1)(Y^2-1)}{16} q^2 v^4 - \frac{(5Y+1)(Y^2-1)}{4} q^2 v^6 \right\}, \quad (\text{B36})$$

$$X_3^{\mathfrak{S}} = 0, \quad (\text{B37})$$

$$X_4^{\mathfrak{S}} = \left\{ -\frac{(Y+1)(Y-1)^2}{32} q^2 v^4 + \frac{(Y+1)(Y-1)^2}{8} q^2 v^6 \right\} \\ +e^2 \left\{ \frac{(Y+1)(Y-1)^2}{32} q^2 v^4 - \frac{(Y+1)(Y-1)^2}{8} q^2 v^6 \right\}, \quad (\text{B38})$$

$$X_n^{\mathfrak{S}} = \begin{cases} 0 & (n: \text{odd}) \\ O(v^{2n-4}) & (n \geq 6: \text{even}) \end{cases}. \quad (\text{B39})$$

C. Secular evolution of the orbital parameters, v , e , and Y

An alternative set of the orbital parameters, $J = \{v, e, Y\}$, is also useful to specify the orbit. The secular changes of the parameters can be derived from those of $I = \{E, L, C\}$, as

$$\left\langle \frac{dJ}{dt} \right\rangle_t = \sum_{I=E,L,C} (G^{-1})_I^J \left\langle \frac{dI}{dt} \right\rangle_t, \quad (\text{C1})$$

where $G_J^I = \partial(E, L, C)/\partial(v, e, Y)$ is the Jacobian matrix for the transformation from $\{E, L, C\}$ to $\{v, e, Y\}$ ³.

Substituting the 3PN $O(e^6)$ formulae of $\langle dI/dt \rangle_t^\infty$ shown in Sec. 3 into the above relation, we obtain the secular changes of $\{v, e, Y\}$ associated with the flux of gravitational waves to infinity as

$$\begin{aligned}
\left\langle \frac{dv}{dt} \right\rangle_t^\infty &= \left(\frac{dv}{dt} \right)_N \left[1 + \frac{7}{8}e^2 + \left\{ -\frac{743}{336} - \frac{55}{21}e^2 + \frac{8539}{2688}e^4 \right\} v^2 \right. \\
&\quad + \left\{ 4\pi - \frac{133}{12}Yq + \left(\frac{97}{8}\pi - \frac{379}{24}Yq \right) e^2 \right. \\
&\quad \left. + \left(\frac{49}{32}\pi - \frac{475}{96}Yq \right) e^4 - \frac{49}{4608}\pi e^6 \right\} v^3 \\
&\quad + \left\{ \frac{34103}{18144} - \frac{329}{96}q^2 + \frac{815}{96}Y^2q^2 + \left(-\frac{526955}{12096} - \frac{929}{96}q^2 + \frac{477}{32}Y^2q^2 \right) e^2 \right. \\
&\quad \left. + \left(-\frac{1232809}{48384} - \frac{1051}{768}q^2 + \frac{999}{256}Y^2q^2 \right) e^4 + \frac{105925}{16128}e^6 \right\} v^4 \\
&\quad + \left\{ -\frac{4159}{672}\pi - \frac{1451}{56}Yq + \left(-\frac{48809}{1344}\pi - \frac{1043}{96}Yq \right) e^2 \right. \\
&\quad + \left(\frac{679957}{43008}\pi - \frac{15623}{336}Yq \right) e^4 + \left(\frac{4005097}{774144}\pi - \frac{35569}{1792}Yq \right) e^6 \left. \right\} v^5 \\
&\quad + \left\{ \frac{16447322263}{139708800} + \frac{16}{3}\pi^2 - \frac{1712}{105}\gamma - \frac{3424}{105}\ln(2) - \frac{331}{192}q^2 \right. \\
&\quad - \frac{289}{6}\pi Yq + \frac{145759}{1344}Y^2q^2 \\
&\quad + \left(\frac{8901670423}{11642400} + \frac{229}{6}\pi^2 - \frac{24503}{210}\gamma + \frac{1391}{30}\ln(2) - \frac{78003}{280}\ln(3) \right. \\
&\quad \left. + \frac{2129}{42}q^2 - \frac{4225}{24}\pi Yq + \frac{27191}{224}Y^2q^2 \right) e^2 \\
&\quad + \left(\frac{269418340489}{372556800} + \frac{109}{4}\pi^2 + \frac{3042117}{1120}\ln(3) - \frac{11663}{140}\gamma \right. \\
&\quad \left. - \frac{418049}{84}\ln(2) - \frac{56239}{10752}q^2 - \frac{17113}{192}\pi Yq + \frac{414439}{3584}Y^2q^2 \right) e^4 \\
&\quad + \left(\frac{174289281}{862400} - \frac{1044921875}{96768}\ln(5) + \frac{23}{16}\pi^2 - \frac{42667641}{3584}\ln(3) \right. \\
&\quad + \frac{94138279}{2160}\ln(2) - \frac{2461}{560}\gamma - \frac{3571}{3584}q^2 - \frac{108577}{13824}\pi Yq \\
&\quad \left. + \frac{41071}{1536}Y^2q^2 \right) e^6
\end{aligned}$$

³ It should be noted that, to calculate the Jacobian matrix up to $O(e^6)$, one need to calculate $\{E, L, C\}$ up to $O(e^8)$ since the leading terms do not depend on e and then the relative orders of accuracy of their first derivatives with the eccentricity of the Jacobian matrix in Eq. (C1) are reduced by $O(e^2)$. For a similar reason, one also need to calculate E up to 5PN order because the relative PN order of $\partial E/\partial v$ is reduced by $O(v^2)$ compared to E .

$$\begin{aligned}
& - \left(\frac{1712}{105} + \frac{24503}{210} e^2 + \frac{11663}{140} e^4 + \frac{2461}{560} e^6 \right) \ln v \Big\} v^6 \Big], \tag{C2} \\
\left\langle \frac{de}{dt} \right\rangle_t^\infty = & \left(\frac{de}{dt} \right)_N \left[1 + \frac{121}{304} e^2 + \left\{ -\frac{6849}{2128} - \frac{2325}{2128} e^2 + \frac{22579}{17024} e^4 \right\} v^2 \right. \\
& + \left\{ \frac{985}{152} \pi - \frac{879}{76} Yq + \left(\frac{5969}{608} \pi - \frac{699}{76} Yq \right) e^2 \right. \\
& \quad \left. + \left(\frac{24217}{29184} \pi - \frac{1313}{608} Yq \right) e^4 \right\} v^3 \\
& + \left\{ -\frac{286397}{38304} - \frac{3179}{608} q^2 + \frac{5869}{608} Y^2 q^2 \right. \\
& \quad + \left(-\frac{2070667}{51072} - \frac{8925}{1216} q^2 + \frac{633}{64} Y^2 q^2 \right) e^2 \\
& \quad \left. + \left(-\frac{3506201}{306432} - \frac{3191}{4864} q^2 + \frac{9009}{4864} Y^2 q^2 \right) e^4 \right\} v^4 \\
& + \left\{ -\frac{1903}{304} Yq - \frac{87947}{4256} \pi + \left(-\frac{3539537}{68096} \pi - \frac{93931}{8512} Yq \right) e^2 \right. \\
& \quad \left. + \left(\frac{5678971}{817152} \pi - \frac{442811}{17024} Yq \right) e^4 \right\} v^5 \\
& + \left\{ -\frac{82283}{1995} \gamma - \frac{11021}{285} \ln(2) - \frac{234009}{5320} \ln(3) + \frac{11224646611}{46569600} + \frac{769}{57} \pi^2 \right. \\
& \quad + \frac{180255}{8512} q^2 - \frac{11809}{152} \pi Yq + \frac{598987}{8512} Y^2 q^2 \\
& \quad + \left(\frac{927800711807}{884822400} - \frac{2982946}{1995} \ln(2) + \frac{2782}{57} \pi^2 + \frac{1638063}{3040} \ln(3) \right. \\
& \quad \left. - \frac{297674}{1995} \gamma + \frac{536653}{8512} q^2 - \frac{91375}{608} \pi Yq + \frac{356845}{8512} Y^2 q^2 \right) e^2 \\
& \quad + \left(\frac{190310746553}{262169600} - \frac{1147147}{15960} \gamma + \frac{10721}{456} \pi^2 - \frac{1022385321}{340480} \ln(3) \right. \\
& \quad + \frac{760314287}{47880} \ln(2) - \frac{1044921875}{204288} \ln(5) + \frac{56509}{9728} q^2 \\
& \quad \left. - \frac{1739605}{29184} \pi Yq + \frac{3248951}{68096} Y^2 q^2 \right) e^4 \\
& \quad \left. - \left(\frac{82283}{1995} + \frac{297674}{1995} e^2 + \frac{1147147}{15960} e^4 \right) \ln v \right\} v^6 \Big], \tag{C3}
\end{aligned}$$

$$\begin{aligned}
\left\langle \frac{dY}{dt} \right\rangle_t^\infty = & \left(\frac{dY}{dt} \right)_N \left[1 + \frac{189}{61} e^2 + \frac{285}{488} e^4 \right. \\
& + \left\{ -\frac{13}{244} Yq - \frac{277}{244} Yq e^2 - \frac{1055}{1952} Yq e^4 \right\} v \\
& + \left\{ -\frac{10461}{1708} - \frac{83723}{3416} e^2 - \frac{21261}{13664} e^4 + \frac{49503}{27328} e^6 \right\} v^2 \\
& + \left\{ \frac{290}{61} \pi - \frac{12755}{3416} Yq + \left(\frac{1990}{61} \pi - \frac{27331}{1708} Yq \right) e^2 \right.
\end{aligned}$$

$$+ \left(\frac{21947}{976} \pi - \frac{540161}{27328} Y q \right) e^4 + \left(\frac{38747}{35136} \pi - \frac{140001}{27328} Y q \right) e^6 \Big\} v^3 \Big], \quad (\text{C4})$$

where the leading contributions are given by

$$\begin{aligned} \left(\frac{dv}{dt} \right)_N &= \frac{32}{5} \left(\frac{\mu}{M^2} \right) v^9 (1 - e^2)^{3/2}, \\ \left(\frac{de}{dt} \right)_N &= -\frac{304}{15} \left(\frac{\mu}{M^2} \right) v^8 e (1 - e^2)^{3/2}, \\ \left(\frac{dY}{dt} \right)_N &= -\frac{244}{15} \left(\frac{\mu}{M^2} \right) v^{11} q (1 - e^2)^{3/2} (1 - Y^2). \end{aligned} \quad (\text{C5})$$

In the same way, substituting the 3.5PN $O(e^6)$ formulae of $\langle dI/dt \rangle_t^H$ shown in Sec. 3 into Eq. (C1), we obtain the secular changes of $\{v, e, Y\}$ associated with the flux of gravitational waves to the horizon as

$$\begin{aligned} \left\langle \frac{dv}{dt} \right\rangle_t^H &= \left(\frac{dv}{dt} \right)_N \left[-\frac{1}{256} \{8 + 24e^2 + 3e^4\} \{8 + 9q^2 + 15Y^2q^2\} Y q v^5 \right. \\ &\quad \left\{ -\frac{11}{8} - \frac{189}{64} q^2 - \frac{15}{64} Y^2 q^2 + \left(-\frac{69}{8} - \frac{81}{4} q^2 + \frac{45}{32} Y^2 q^2 \right) e^2 \right. \\ &\quad \left. + \left(-\frac{381}{64} - \frac{7479}{512} q^2 + \frac{1035}{512} Y^2 q^2 \right) e^4 \right. \\ &\quad \left. + \left(-\frac{11}{32} - \frac{423}{512} q^2 + \frac{45}{512} Y^2 q^2 \right) e^6 \right\} Y q v^7 \Big], \end{aligned} \quad (\text{C6})$$

$$\begin{aligned} \left\langle \frac{de}{dt} \right\rangle_t^H &= \left(\frac{de}{dt} \right)_N \left[-\frac{33}{4864} \{8 + 12e^2 + e^4\} \{8 + 9q^2 + 15Y^2q^2\} Y q v^5 \right. \\ &\quad \left\{ -\frac{453}{152} - \frac{8127}{1216} q^2 - \frac{45}{1216} Y^2 q^2 + \left(-\frac{2979}{304} - \frac{28593}{1216} q^2 + \frac{1485}{608} Y^2 q^2 \right) e^2 \right. \\ &\quad \left. + \left(-\frac{5649}{1216} - \frac{111591}{9728} q^2 + \frac{16515}{9728} Y^2 q^2 \right) e^4 \right\} Y q v^7 \Big], \end{aligned} \quad (\text{C7})$$

$$\begin{aligned} \left\langle \frac{dY}{dt} \right\rangle_t^H &= \left(\frac{dY}{dt} \right)_N \left[-\frac{3}{7808} \{8 + 24e^2 + 3e^4\} \{16 + 33q^2 + 15Y^2q^2\} v^2 \right. \\ &\quad \left\{ -\frac{51}{122} + \frac{585}{1952} Y^2 q^2 - \frac{1953}{1952} q^2 + \left(-\frac{225}{61} - \frac{16875}{1952} q^2 + \frac{3375}{1952} Y^2 q^2 \right) e^2 \right. \\ &\quad \left. + \left(-\frac{2961}{976} - \frac{109863}{15616} q^2 + \frac{16335}{15616} Y^2 q^2 \right) e^4 \right. \\ &\quad \left. + \left(-\frac{171}{976} - \frac{3159}{7808} q^2 + \frac{405}{7808} Y^2 q^2 \right) e^6 \right\} v^4 \Big]. \end{aligned} \quad (\text{C8})$$

Actually, we can obtain the higher PN results by using the 4PN $O(e^6)$ formulae for the secular changes of $\{E, L, C\}$, although we do not present them in the text. The full expressions of $\langle dJ/dt \rangle_t^\infty$ and $\langle dJ/dt \rangle_t^H$ for $J = \{v, e, Y\}$ will be available online [26].

Here we make a comment on the reliable order of the expansion with respect to e in $\langle de/dt \rangle_t$. By using Eq. (C1), $\langle de/dt \rangle_t$ can be calculated from the linear combination of the secular changes of $\{E, L, C\}$. Since the leading order of the (e, I) -component of the inverse Jacobian matrix is $O(1/e)$, each term in the linear combination is apparently $O(1/e)$.




Fig. C1 The relative errors in the analytic formulae for the secular change of the orbital eccentricity due to the gravitational waves to infinity. We plot Δ_e , defined in a similar manner to Eq. (39), as a function of the semi-latus rectum p for $q = 0.9$, $e = 0.1, 0.4$ and 0.7 (from left to right) and $\iota = 50^\circ$. We truncated the plots at $p = 6$ because the relative errors get too large in $p < 6$ to be meaningful. The relative error in the previous 2.5PN $O(e^2)$ formula given in [15] strays off the p^{-3} line earlier than the 2.5PN $O(e^2)$ formula in this paper. This trend is clearer for larger e . The relative errors in the 3PN $O(e^4)$ and 4PN $O(e^4)$ formulae fall off faster than p^{-3} and p^{-4} for small e cases as expected, while this is not the case for $e = 0.7$ because of the higher order correction of e than $O(e^4)$.

However, the $O(1/e)$ contribution turns out to vanish due to a cancellation in taking the combination, and hence $\langle de/dt \rangle_t$ is $O(e)$, which corresponds to the well-known fact that circular orbits remain circular [40, 41], *i.e.* $\langle de/dt \rangle_t = 0$ when $e = 0$. This cancellation reduces the reliable order in $\langle de/dt \rangle_t$ by $O(e^2)$, compared to the order of $\langle dI/dt \rangle_t$ for $I = \{E, L, C\}$. Since we calculate $\langle dI/dt \rangle_t$ up to $O(e^6)$ in this paper, we can obtain $\langle de/dt \rangle_t$ correctly up to $O(e^4)$ from the leading order.

$\langle dv/dt \rangle_t^\infty$ and $\langle dY/dt \rangle_t^\infty$ in Eqs. (C2) and (C4) are consistent up to the 2.5PN $O(e^2)$ order with the previous results in Ref. [15], while we find inconsistency in the $O(e^2)$ terms of the formula for $\langle de/dt \rangle_t^\infty$ in [15]. This may be explained by the reduction in the reliable order mentioned above: the calculations of $\langle dI/dt \rangle_t$ in [15] are done up to $O(e^2)$, and therefore the resultant formula of $\langle de/dt \rangle_t$ is reliable only at the leading order. We can also confirm it numerically. In Fig. C1 we show the relative errors in the two analytic formulae by comparing to numerical results [18] in a similar manner to Eq. (39). It can be found that the relative error in the previous 2.5PN $O(e^2)$ formula strays out of the expected power law, p^{-3} , earlier than that in our 2.5PN $O(e^2)$ formula. This trend is clearer for larger eccentricity. We can also confirm the validity of our formula by seeing that the relative errors in our 4PN $O(e^4)$ formula falls off faster than p^{-4} . (The leading PN order of the difference in the $O(e^2)$ terms between the previous and our formulae is $\frac{39}{38}e^2v^2$. If our formula contains any error in the e^2v^2 term, the relative error will not fall off faster than p^{-4} for large p .)

A similar reduction in the PN order occurs in the calculation of $\langle dY/dt \rangle_t^\infty$: although each term in the linear combination of Eq. (C1) for $J = Y$ is $O(v^8)$, the terms at the first two orders, $O(v^8)$ and $O(v^{10})$, vanish due to a cancellation in taking the combination. As a result, the leading order of $\langle dY/dt \rangle_t$ is $O(v^{11})$ and hence the reliable order relative to the leading term is reduced to $O(v^5)$ (2.5PN order) when we have $\langle dI/dt \rangle_t$ for $I = \{E, L, C\}$ up to $O(v^8)$ (4PN order).




Fig. C2 The relative errors in the 4PN formulae for the secular changes of the orbital parameters, $\{v, e, Y\}$. We plot the relative errors, Δ_J defined in a similar manner to Eq. (39), as functions of the semi-latus rectum p for $q = 0.9$, $e = 0.1, 0.4$ and 0.7 (from left to right) and $\iota = 50^\circ$. We truncated the plots at $p = 6$ because the relative errors get too large in $p < 6$ to be meaningful. Δ_v and Δ_e fall off faster than p^{-4} , while Δ_Y approximately fall off as p^{-3} , slower than $O(p^{-4})$. This confirms that the relative order of the PN correction of the analytic formula for $\langle dY/dt \rangle_t$ is reduced from 4PN to 2.5PN because of the cancellation of the low PN terms.

In Fig. C2, we show the relative errors in the analytic PN formulae for the secular changes of the orbital parameters, $\{v, e, Y\}$, derived from the 4PN $O(e^6)$ formulae of $\langle dI/dt \rangle_t$ for $I = \{E, L, C\}$. Similarly in Fig. 1, the relative errors in the analytic formulae for $\langle dv/dt \rangle_t$ and $\langle de/dt \rangle_t$ as functions of the semi-latus rectum p fall off faster than p^{-4} when the eccentricity is small. Observe, however, that the relative error in the analytic formula for $\langle dY/dt \rangle_t$ falls off faster than $p^{-5/2}$, but slower than p^{-4} .

From the leading order expressions in Eq. (C5), one will find the well known fact that equatorial orbits stay in the equatorial plane [10, 15, 42], *i.e.* $\langle dY/dt \rangle_t = 0$ when $Y = 1$. In the Schwarzschild case ($q = 0$), the secular changes of v and e do not depend on Y in addition to $\langle dY/dt \rangle_t = 0$. This implies that the orbital plane can be fixed on the equatorial plane ($\theta = \pi/2$) due to the spherical symmetry of Schwarzschild spacetime.

One will also find that the radiation reaction reduces the orbital eccentricity and increases the orbital velocity since $(de/dt)_N \leq 0$ and $(dv/dt)_N \geq 0$ [40, 41], while the radiation reaction increases (decreases) the inclination angle since $(dY/dt)_N \leq 0$ ($(dY/dt)_N \geq 0$) when $q \geq 0$ ($q \leq 0$) [10, 15, 42]. Moreover, the secular change of the inclination angle is smaller than those of the other orbital parameters since $\langle d \ln e / d \ln v \rangle_t = O(v^0)$ and $\langle d \ln Y / d \ln v \rangle_t = O(v^3)$.

References

- [1] Y. Mino, M. Sasaki and T. Tanaka, Phys. Rev. D **55** (1997) 3457 [gr-qc/9606018].
- [2] T. C. Quinn and R. M. Wald, Phys. Rev. D **56** (1997) 3381 [gr-qc/9610053].
- [3] E. Poisson, A. Pound and I. Vega, Living Rev. Rel. **14** (2011) 7 [arXiv:1102.0529 [gr-qc]].
- [4] L. Barack, Class. Quant. Grav. **26** (2009) 213001 [arXiv:0908.1664 [gr-qc]].
- [5] B. Wardell, arXiv:1501.07322 [gr-qc].
- [6] T. Hinderer and E. E. Flanagan, Phys. Rev. D **78** (2008) 064028 [arXiv:0805.3337 [gr-qc]].
- [7] E. E. Flanagan and T. Hinderer, Phys. Rev. Lett. **109** (2012) 071102 [arXiv:1009.4923 [gr-qc]].
- [8] Y. Mino, Phys. Rev. D **67** (2003) 084027 [gr-qc/0302075].
- [9] N. Sago, T. Tanaka, W. Hikida and H. Nakano, Prog. Theor. Phys. **114** (2005) 509 [gr-qc/0506092].
- [10] N. Sago, T. Tanaka, W. Hikida, K. Ganz and H. Nakano, Prog. Theor. Phys. **115** (2006) 873 [gr-qc/0511151].
- [11] T. Osburn, E. Forseth, C. R. Evans and S. Hopper, Phys. Rev. D **90** (2014) 10, 104031 [arXiv:1409.4419 [gr-qc]].

-
- [12] Y. Mino, M. Sasaki, M. Shibata, H. Tagoshi and T. Tanaka, Prog. Theor. Phys. Suppl. **128** (1997) 1 [gr-qc/9712057].
 - [13] R. Fujita, Prog. Theor. Phys. **128** (2012) 971 [arXiv:1211.5535 [gr-qc]].
 - [14] R. Fujita, Prog. Theor. Exp. Phys. **2015** 033E01 (2015) [arXiv:1412.5689 [gr-qc]].
 - [15] K. Ganz, W. Hikida, H. Nakano, N. Sago and T. Tanaka, Prog. Theor. Phys. **117** (2007) 1041 [gr-qc/0702054].
 - [16] R. Fujita and H. Tagoshi, Prog. Theor. Phys. **112** (2004) 415 [gr-qc/0410018].
 - [17] R. Fujita and H. Tagoshi, Prog. Theor. Phys. **113** (2005) 1165 [arXiv:0904.3818 [gr-qc]].
 - [18] R. Fujita, W. Hikida and H. Tagoshi, Prog. Theor. Phys. **121** (2009) 843 [arXiv:0904.3810 [gr-qc]].
 - [19] S. Drasco and S. A. Hughes, Phys. Rev. D **69** (2004) 044015 [astro-ph/0308479].
 - [20] S. Isoyama, R. Fujita, H. Nakano, N. Sago and T. Tanaka, PTEP **2013** (2013) 6, 063E01 [arXiv:1302.4035 [gr-qc]].
 - [21] S. A. Teukolsky, Astrophys. J. **185** (1973) 635.
 - [22] M. Sasaki and H. Tagoshi, Living Rev. Rel. **6** (2003) 6 [gr-qc/0306120].
 - [23] S. A. Teukolsky and W. H. Press, Astrophys. J. **193** (1974) 443.
 - [24] S. Drasco, E. E. Flanagan and S. A. Hughes, Class. Quant. Grav. **22** (2005) S801 [gr-qc/0505075].
 - [25] D.V. Gal'tsov, J. Phys. A **15**, 3737 (1982).
 - [26] Kyoto University, Japan. <http://www2.yukawa.kyoto-u.ac.jp/~misao.sasaki/BHPC/index.html>, date last accessed April 8, 2015.
 - [27] E. Poisson and M. Sasaki, Phys. Rev. D **51** (1995) 5753 [gr-qc/9412027].
 - [28] H. Tagoshi, S. Mano and E. Takasugi, Prog. Theor. Phys. **98** (1997) 829 [gr-qc/9711072].
 - [29] S. A. Hughes, Phys. Rev. D **64** (2001) 064004 [gr-qc/0104041].
 - [30] D. V. Galtsov, A. A. Matyukhin and V. I. Petukhov, Phys. Lett. A **77**, 387-390 (1980)
 - [31] M. Shibata, Phys. Rev. D **50**, 6297-6311 (1994)
 - [32] K. G. Arun, L. Blanchet, B. R. Iyer and M. S. S. Qusailah, Phys. Rev. D **77**, 064035 (2008) [arXiv:0711.0302 [gr-qc]].
 - [33] K. G. Arun, L. Blanchet, B. R. Iyer and S. Sinha, Phys. Rev. D **80**, 124018 (2009) [arXiv:0908.3854 [gr-qc]].
 - [34] T. Damour, B. R. Iyer and B. S. Sathyaprakash, Phys. Rev. D **57** (1998) 885 [gr-qc/9708034].
 - [35] T. Damour and A. Nagar, Phys. Rev. D **76** (2007) 064028 [arXiv:0705.2519 [gr-qc]].
 - [36] T. Damour and A. Nagar, Phys. Rev. D **77** (2008) 024043 [arXiv:0711.2628 [gr-qc]].
 - [37] T. Damour, B. R. Iyer and A. Nagar, Phys. Rev. D **79** (2009) 064004 [arXiv:0811.2069 [gr-qc]].
 - [38] S. Isoyama, R. Fujita, N. Sago, H. Tagoshi and T. Tanaka, Phys. Rev. D **87** (2013) 2, 024010 [arXiv:1210.2569 [gr-qc]].
 - [39] R. Fujita and W. Hikida, Class. Quant. Grav. **26** (2009) 135002 [arXiv:0906.1420 [gr-qc]].
 - [40] P. C. Peters, Phys. Rev. **136**, 1224-1232 (1964).
 - [41] H. Tagoshi, Prog. Theor. Phys. **93** 307-333 (1995).
 - [42] M. Shibata, M. Sasaki, H. Tagoshi and T. Tanaka, Phys. Rev. D **51** (1995) 1646-1663 [gr-qc/9409054].